

ON THE TRANSITION TO THE NORMAL PHASE FOR SUPERCONDUCTORS SURROUNDED BY NORMAL CONDUCTORS

SØREN FOURNAIS AND AYMAN KACHMAR

ABSTRACT. For a cylindrical superconductor surrounded by a normal material, we discuss transition to the normal phase of stable, locally stable and critical configurations. Associated with those phase transitions, we define critical magnetic fields and we provide a sufficient condition for which those critical fields coincide. In particular, when the conductivity ratio of the superconducting and the normal material is large, we show that the aforementioned critical magnetic fields coincide, thereby proving that the transition to the normal phase is sharp. One key-ingredient in the paper is the analysis of an elliptic boundary value problem involving ‘transmission’ boundary conditions. Another key-ingredient involves a monotonicity result (with respect to the magnetic field strength) of the first eigenvalue of a magnetic Schrödinger operator with discontinuous coefficients.

CONTENTS

1. Introduction	1
2. Proof of Theorem 1.1	5
2.1. Basic estimates on solutions	5
2.2. Weak Decay Estimate and applications	10
3. Monotonicity of the first eigenvalue	13
3.1. The constant $\alpha_0(a, m)$	13
3.2. Notation, hypotheses and announcement of Main Result	16
3.3. Some facts concerning $\mu_1(B, \alpha)$	20
3.4. Proof of Theorem 3.3: General domains	20
3.5. Proof of Theorem 3.4: Disc domains	24
4. Exponential decay of order parameters	24
Acknowledgements	26
Appendix A. Improved eigenvalue estimate for the disc	26
References	30

1. INTRODUCTION

A type II superconductor placed in an applied magnetic field undergoes three phase transitions: When the intensity of the applied field is below a first critical value H_{C_1} , the sample exhibits the Meissner effect and remains in a superconducting phase. When the field is increased further past H_{C_1} , the sample is in a mixed state and the magnetic field penetrates the material in quantized vortices. Increasing

2000 *Mathematics Subject Classification.* Primary 35Q55, Secondary 35J25, 35A15, 58E50, 81Q10.

Key words and phrases. Ginzburg-Landau type equations, Schrödinger operator with magnetic field, semiclassical analysis.

the field further past another critical value H_{C_2} , the sample exhibits surface superconductivity, and when the field is increased above H_{C_3} , superconductivity is lost and the sample is in the normal phase. The above picture is rigorously established for extreme type II materials through the minimization of the Ginzburg-Landau functional, see for instance the papers [8, 12, 20, 27, 28, 30] and the books [7, 29] for results and additional references concerning the subject.

In addition to the phase transitions associated with minimizers (stable states) of the Ginzburg-Landau functional, type II materials possess hysteresis associated with local minimizers (locally stable states) of the energy. For instance, a locally stable state that does not possess vortices will remain locally stable in increasing magnetic fields up to a *super-heating field*, and a similar phenomenon is associated with a *sub-cooling field* associated with decreasing applied fields (see [31]). It is therefore natural to address a similar question when dealing with the transition of normal states: Does local stability persist for the normal state below H_{C_3} , and for the superconducting state above H_{C_3} , or will there be hysteresis? That is, we ask whether the transition from the superconducting phase to the normal phase happens at a sharp critical value of the magnetic field ($= H_{C_3}$).

For type II superconducting samples with smooth boundaries and surrounded by the vacuum, Fournais and Helffer [8] showed that the transition is indeed sharp. Hysteresis is excluded in [7]. The case of domains with corners [3] and the 3-dimensional case [8, 26] (see also [19, 26]) have also been studied. The reason for the sharp transition is essentially the monotonicity of the first eigenvalue $\mu_1(B)$ of the Neumann Schrödinger operator

$$-(\nabla - iB\mathbf{F})^2 \quad \text{in } \Omega, \quad \Omega \subset \mathbb{R}^d \quad (d = 2, 3),$$

for large values of the magnetic field, a property known as strong diamagnetism (see [8, 9]). Here $\mathbf{F} : \bar{\Omega} \rightarrow \mathbb{R}^d$ is a vector field such that $\text{curl } \mathbf{F}$ is a constant.

In this paper, we address the same question—the transition from superconducting to normal phase—but for a superconductor surrounded by normal materials. It is well known from the superconducting proximity effect (see [13, 14]) that the presence of a normal material exterior to a superconductor allows the superconducting electron Cooper pairs to flow into the normal material in a narrow boundary layer. The characteristic length scale of that layer is called the ‘extrapolation length’. To model this phenomenon, one has to consider a generalized Ginzburg-Landau theory where the order parameter and the magnetic potential are not only defined in the superconducting material but also in the normal material surrounding it.

For a cylindrical superconducting sample of cross section $\Omega \subset \mathbb{R}^2$, surrounded by a normal material and placed in a magnetic field parallel to the cylinder axis, the Gibbs free energy is given by the following Ginzburg-Landau type functional (see [4]):

$$\begin{aligned} \mathcal{G}(\psi, \mathbf{A}) = \mathcal{G}_{\kappa, H}(\psi, \mathbf{A}) = & \int_{\Omega} \left(|\nabla_{\kappa H \mathbf{A}} \psi|^2 - \kappa^2 |\psi|^2 + \frac{\kappa^2}{2} |\psi|^4 \right) dx \\ & + \int_{\tilde{\Omega} \setminus \Omega} \left(\frac{1}{m} |\nabla_{\kappa H \mathbf{A}} \psi|^2 + a \kappa^2 |\psi|^2 \right) dx \\ & + (\kappa H)^2 \int_{\tilde{\Omega}} |\text{curl } \mathbf{A} - 1|^2 dx. \end{aligned} \quad (1.1)$$

Here we use the notation,

$$\nabla_{\mathbf{A}} = \nabla - i\mathbf{A}, \quad (1.2)$$

for the magnetic gradient.

In the above functional, $\tilde{\Omega} \subset \mathbb{R}^2$ is a bounded domain such that $\bar{\Omega} \subset \tilde{\Omega}$ and $\tilde{\Omega} \setminus \Omega$ is the cross section of the normal material¹, the Ginzburg-Landau parameter $\kappa > 0$ is a characteristic of the superconducting material (the ratio of two characteristic lengths), $H > 0$ is the intensity of the applied magnetic field, $a > 0$ is a characteristic of the normal material that depends on the temperature and its sign signifies that the temperature is above the critical temperature of the normal material. Finally, $m > 0$ is the conductivity ratio of the superconducting and normal materials. Minimization of the functional (1.1) will take place over finite-energy configurations $(\psi, \mathbf{A}) \in H^1(\tilde{\Omega}; \mathbb{C}) \times H^1(\tilde{\Omega}; \mathbb{R}^2)$. Starting from a minimizing sequence, it is quite standard to prove the existence of minimizers of (1.1), see [16].

We will always assume that Ω and $\tilde{\Omega}$ are smooth, bounded and simply connected.

Notice that the normal state $(0, \mathbf{F})$, with $\mathbf{F}$ being the unique vector field satisfying

$$\operatorname{curl} \mathbf{F} = 1, \quad \operatorname{div} \mathbf{F} = 0 \quad \text{in } \tilde{\Omega}, \quad \nu \cdot \nabla \mathbf{F} = 0 \quad \text{on } \partial \tilde{\Omega}, \quad (1.3)$$

is a critical point of the functional (1.1). It can also be shown that this state is the unique normal state up to a *gauge transformation* (see [16]). Configurations which are gauge equivalent to the normal state $(0, \mathbf{F})$ will be called *trivial* throughout the paper.

Defining the set,

$$\mathcal{N}^{\text{sc}}(a, m; \kappa) = \{H > 0 : \mathcal{G}_{\kappa, H} \text{ has non-trivial critical points}\}, \quad (1.4)$$

then it is known from [16, Theorem 4.7] (see also Theorem 2.3 in the present paper) that the above set is bounded.

In connection with stability and local stability of the normal state $(0, \mathbf{F})$, we also introduce the two sets:

$$\begin{aligned} \mathcal{N}(a, m; \kappa) &= \{H > 0 : \mathcal{G}_{\kappa, H} \text{ has a non-trivial minimizer}\}, \\ \mathcal{N}^{\text{loc}}(a, m; \kappa) &= \{H > 0 : \mu^{(1)}(\kappa, H) < 0\}. \end{aligned} \quad (1.5)$$

Here $\mu^{(1)}(\kappa, H)$ is the ground state energy of the quadratic form

$$\mathcal{Q}[\kappa, H](\phi) = \int_{\Omega} (|\nabla_{\kappa H \mathbf{F}} \phi|^2 - \kappa^2 |\phi|^2) dx + \int_{\tilde{\Omega} \setminus \Omega} \left(\frac{1}{m} |\nabla_{\kappa H \mathbf{F}} \phi|^2 + a \kappa^2 |\phi|^2 \right) dx, \quad (1.6)$$

i.e.

$$\mu^{(1)}(\kappa, H) = \inf_{\substack{\phi \in H^1(\tilde{\Omega}) \\ \phi \neq 0}} \left(\frac{\mathcal{Q}[\kappa, H](\phi)}{\|\phi\|_{L^2(\tilde{\Omega})}^2} \right). \quad (1.7)$$

Since $\mathcal{Q}[\kappa, H]$ defines the Hessian of the functional $\mathcal{G}_{\kappa, H}$ at the normal state $(0, \mathbf{F})$, we see that if $H \in \mathcal{N}^{\text{loc}}(a, m; \kappa)$, then $(0, \mathbf{F})$ is not a local minimizer $\mathcal{G}_{\kappa, H}$. Hence we obtain the following inclusion,

$$\mathcal{N}^{\text{loc}}(a, m; \kappa) \subset \mathcal{N}(a, m; \kappa).$$

On the other hand, the following inclusion is trivial,

$$\mathcal{N}(a, m; \kappa) \subset \mathcal{N}^{\text{sc}}(a, m; \kappa). \quad (1.8)$$

One of the main results of the present paper is the following.

¹In part of the existing literature on the subject (see for example [23, 4]) $\tilde{\Omega}$ is taken to be all of $\mathbb{R}^2$. However, in the original paper of [15], the normal material (i.e. $\tilde{\Omega} \setminus \Omega$) has been taken to be bounded. We take the latter point of view, in order to avoid certain technical difficulties which are unimportant for our present purpose. The functional analysis necessary to overcome those difficulties is developed in [16].

Theorem 1.1. *Given $a > 0$ and $m > 1$, there exists $\kappa_0 > 0$ such that, for all $\kappa \geq \kappa_0$, the following equalities hold,*

$$\mathcal{N}^{\text{sc}}(a, m; \kappa) = \mathcal{N}^{\text{loc}}(a, m; \kappa) = \mathcal{N}(a, m; \kappa).$$

In the literature, it is typical to introduce the following critical fields (cf. e.g. [8, 27]),

$$\overline{H}_{C_3}^{\text{sc}}(a, m; \kappa) = \sup \mathcal{N}^{\text{sc}}(a, m; \kappa), \quad \underline{H}_{C_3}^{\text{sc}}(a, m; \kappa) = \inf \mathbb{R}_+ \setminus \mathcal{N}^{\text{sc}}(a, m; \kappa), \quad (1.9)$$

$$\overline{H}_{C_3}(a, m; \kappa) = \sup \mathcal{N}(a, m; \kappa), \quad \underline{H}_{C_3}(a, m; \kappa) = \inf \mathbb{R}_+ \setminus \mathcal{N}(a, m; \kappa), \quad (1.10)$$

$$\overline{H}_{C_3}^{\text{loc}}(a, m; \kappa) = \sup \mathcal{N}^{\text{loc}}(a, m; \kappa), \quad \underline{H}_{C_3}^{\text{loc}}(a, m; \kappa) = \inf \mathbb{R}_+ \setminus \mathcal{N}^{\text{loc}}(a, m; \kappa), \quad (1.11)$$

As a corollary of Theorem 1.1, we get a result concerning equality of the above critical fields.

Corollary 1.2. *Given $a > 0$ and $m > 1$, there exists $\kappa_0 > 0$ such that, for all $\kappa \geq \kappa_0$, the following equalities hold,*

$$\overline{H}_{C_3}^{\text{sc}}(a, m; \kappa) = \overline{H}_{C_3}(a, m; \kappa) = \overline{H}_{C_3}^{\text{loc}}(a, m; \kappa),$$

and

$$\underline{H}_{C_3}^{\text{sc}}(a, m; \kappa) = \underline{H}_{C_3}(a, m; \kappa) = \underline{H}_{C_3}^{\text{loc}}(a, m; \kappa).$$

Remark 1.3. *In [23], the second author has established the following asymptotic expansion of $\underline{H}_{C_3}^{\text{loc}}(a, m; \kappa)$ (it can also be shown to hold for $\overline{H}_{C_3}^{\text{loc}}(a, m; \kappa)$):*

$$\underline{H}_{C_3}^{\text{loc}}(a, m; \kappa) = \frac{\kappa}{\alpha_0(a, m)} (1 + o(1)) \quad \text{as } \kappa \rightarrow \infty. \quad (1.12)$$

The constant $\alpha_0(a, m)$ in (1.12) satisfies $\frac{1}{2} < \alpha_0(a, m) < 1$ for $m > 1$ and $\alpha_0(a, m) = 1$ otherwise, and is defined via an auxiliary spectral problem (see Theorem 3.1 below).

Remark 1.4. *Theorem 1.1 does not cover the regime $m \leq 1$. In this specific regime, we have by Remark 1.3 that the nucleation field H_{C_3} coincides with the second critical field H_{C_2} . This reflects one feature of superconductors surrounded by normal materials, that surface superconductivity can be absent (see [15]). However, we need specific tools to treat this interesting case, which are beyond the scope of the present paper.*

We say that the transition from the superconducting to the normal phase is sharp if we have equality between upper and lower fields. By Corollary 1.2 it suffices to verify this for the ‘local fields’, i.e. look whether the following equality holds:

$$\underline{H}_{C_3}^{\text{loc}}(a, m; \kappa) = \overline{H}_{C_3}^{\text{loc}}(a, m; \kappa)?$$

One part of the present paper is devoted to this question, which links to a monotonicity problem of magnetic Schrödinger operators.

It is shown in [23] that there exists $m_* \geq 1$ (see (3.15) below for a precise definition of m_*) such that for $m > m_*$, the second correction term in (1.12) is of order 1 and determined by the maximal value of the scalar curvature of Ω . The precise result in this case is the following:

$$\underline{H}_{C_3}(a, m; \kappa) = \frac{\kappa}{\alpha_0(a, m)} + \frac{\mathcal{C}_1(a, m)}{\alpha_0(a, m)^{3/2}} (\kappa_r)_{\max} + \mathcal{O}(\kappa^{-1/3}), \quad \text{as } \kappa \rightarrow +\infty, \quad (1.13)$$

where the function $\mathcal{C}_1(a, \cdot) : [m_*, +\infty[\mapsto \mathbb{R}_+$ is defined via an auxiliary self-adjoint operator (see (3.14)) and κ_r denotes the scalar curvature of $\partial\Omega$.

Our final result is that all the critical fields coincide for large κ under the extra condition that m is sufficiently large.

Theorem 1.5. *Given $\Omega, \tilde{\Omega}$ and $a > 0$, there exists $m_0 > 0$ and if $m > m_0$, there exists $\kappa_0 > 0$ such that if $\kappa > \kappa_0$ then all the six critical fields defined in (1.9), (1.10) and (1.11) coincide. Furthermore, their common value is the unique solution $H = H_{C_3}(a, m; \kappa)$ of the equation*

$$\mu^{(1)}(\kappa, H) = 0.$$

In particular, the asymptotics (1.13) holds for all the six different definitions of the critical field.

As we mentioned earlier, the essential key for establishing Theorem 1.5 is a monotonicity result concerning the first eigenvalue of a Schrödinger type operator. Actually, let $\mu_1(B, \alpha)$ be the first eigenvalue of the following operator

$$-\nabla_{B\mathbf{F}} \cdot w_m(x) \nabla_{B\mathbf{F}} + \alpha B V_a(x) \quad \text{in } \tilde{\Omega}, \quad (1.14)$$

with

$$w_m(x) = \begin{cases} 1 & \text{in } \Omega \\ \frac{1}{m} & \text{in } \mathbb{R}^2 \setminus \Omega, \end{cases} \quad V_a(x) = \begin{cases} -1 & \text{in } \Omega \\ a & \text{in } \mathbb{R}^2 \setminus \Omega. \end{cases} \quad (1.15)$$

Roughly speaking, we will establish that the equation in (B, α)

$$\mu_1(B, \alpha) = 0$$

admits a unique solution provided that B is large enough and α remains close to $\alpha_0(a, m)$.

In order to prove Theorem 1.1, we establish the following crucial estimate, valid for any critical point of the functional $\mathcal{G}_{\kappa, H}$ (i.e. solution of (2.1) below),

$$a \int_{\tilde{\Omega} \setminus \Omega} |\psi|^4 dx \leq \int_{\Omega} |\psi|^4 dx.$$

The above estimate is non-trivial and we prove it through a detailed analysis of the regularity of critical points of the functional (1.1) (see Lemma 2.5 and Theorem 2.4 below). With the above estimate on the one hand, and other weak decay estimates established in Lemmas 2.2 and 2.6 on the other hand, we prove Theorem 2.7, which links the existence of non-trivial critical points to a spectral condition on the eigenvalue (1.7). Theorem 1.1 is then a consequence of Theorem 2.7.

The paper is organized in the following way. In Section 2, we establish a necessary and sufficient condition for the functional $\mathcal{G}_{\kappa, H}$ to admit non-trivial critical points, see Theorem 2.7.

In Section 3 we establish a monotonicity result for the eigenvalue $\mu_1(B, \alpha)$, see Theorem 3.3. As a consequence, we obtain equality of local fields stated in Theorem 1.5, Theorem 3.5 and Remark 3.6.

In Section 4, we discuss the decay of energy minimizing order parameters, and we prove that they decay exponentially away from the boundary provided that the magnetic field is sufficiently large.

Finally, in the appendix, we prove an improved expansion for (1.13) in the particular case when Ω is a disc domain.

2. PROOF OF THEOREM 1.1

2.1. Basic estimates on solutions. We return now to the analysis of critical points of the functional (1.1). As we mentioned in the introduction, minimizers of

(1.1) exist in the space $H^1(\tilde{\Omega}; \mathbb{C}) \times H^1(\tilde{\Omega}; \mathbb{R}^2)$ and they are *weak* solutions of the associated Ginzburg-Landau equations:

$$\begin{cases} -\nabla_{\kappa H \mathbf{A}} \cdot w_m \nabla_{\kappa H \mathbf{A}} \psi + \kappa^2 (V_a \psi + \mathbf{1}_\Omega |\psi|^2 \psi) = 0, \\ \nabla^\perp \text{curl } \mathbf{A} = (\kappa H)^{-1} w_m \text{Im}(\bar{\psi} (\nabla - i\kappa H \mathbf{A}) \psi), \quad \text{in } \tilde{\Omega}, \\ \nu \cdot (\nabla - i\kappa H \mathbf{A}) \psi = 0, \quad \text{curl } \mathbf{A} = 1 \quad \text{on } \partial \tilde{\Omega}, \end{cases} \quad (2.1)$$

where ν is the unit outward normal vector of $\partial \tilde{\Omega}$.

In order to avoid the question of regularity concerning solutions of the above equation, we shall invoke energy arguments and use only the weak formulation of (2.1). More precisely, $(\psi, \mathbf{A}) \in H^1(\tilde{\Omega}; \mathbb{C}) \times H^1(\tilde{\Omega}; \mathbb{R}^2)$ is a weak solution of (2.1) if for all $\phi \in H^1(\tilde{\Omega}; \mathbb{C})$ and $\mathbf{a} \in H^1(\tilde{\Omega}; \mathbb{R}^2)$, the following equalities hold,

$$\begin{aligned} \int_{\tilde{\Omega}} \left(w_m(x) (\nabla - i\kappa H \mathbf{A}) \psi \cdot \overline{(\nabla - i\kappa H \mathbf{A}) \phi} \right. \\ \left. + \kappa^2 (V_a(x) \psi + \mathbf{1}_\Omega(x) |\psi|^2 \psi) \bar{\phi} \right) dx = 0, \end{aligned} \quad (2.2)$$

$$\int_{\tilde{\Omega}} \text{curl}(\mathbf{A} - \mathbf{F})(\text{curl } \mathbf{a}) - w_m(x) (\kappa H)^{-1} \text{Im}(\bar{\psi} (\nabla - i\kappa H \mathbf{A}) \psi) \cdot \mathbf{a} dx = 0. \quad (2.3)$$

A standard choice of gauge permits us to minimize (1.1) in the reduced space $H^1(\tilde{\Omega}; \mathbb{C}) \times H_{\text{div}}^1(\tilde{\Omega}; \mathbb{R}^2)$, where

$$H_{\text{div}}^1(\tilde{\Omega}; \mathbb{R}^2) = \{ \mathbf{A} \in H^1(\tilde{\Omega}; \mathbb{R}^2) : \text{div } \mathbf{A} = 0 \quad \text{in } \tilde{\Omega}, \quad \nu \cdot \nabla \mathbf{A} = 0 \quad \text{on } \partial \tilde{\Omega} \}. \quad (2.4)$$

We get as immediate consequence of the Poincaré Lemma (see e.g. [32, p. 16, Theorem 1.5]):

Lemma 2.1. *There exists a constant $C > 0$ such that for all $\mathbf{A} \in H_{\text{div}}^1(\tilde{\Omega}; \mathbb{R}^2)$,*

$$\| \mathbf{A} - \mathbf{F} \|_{H^1(\tilde{\Omega})} \leq C \| \text{curl } \mathbf{A} - 1 \|_{L^2(\tilde{\Omega})}. \quad (2.5)$$

Here $\mathbf{F}$ is the vector field introduced in (1.3).

Lemma 2.2. *Let $(\psi, \mathbf{A})$ be a critical configuration of (1.1), i.e. a weak solution of (2.1). Then the following estimates hold :*

$$\| \psi \|_{L^\infty(\tilde{\Omega})} \leq 1, \quad (2.6)$$

$$\| (\nabla - i\kappa H \mathbf{A}) \psi \|_{L^2(\tilde{\Omega})} \leq \max(1, \sqrt{m}) \kappa \| \psi \|_{L^2(\Omega)}, \quad (2.7)$$

$$a \int_{\tilde{\Omega} \setminus \Omega} |\psi|^2 dx \leq \int_{\Omega} |\psi|^2 dx, \quad (2.8)$$

$$\| \psi \|_{L^4(\Omega)}^2 \leq \| \psi \|_{L^2(\Omega)}, \quad (2.9)$$

and

$$H \| \text{curl } \mathbf{A} - 1 \|_{L^2(\tilde{\Omega})} \leq C \| \psi \|_{L^2(\Omega)} \| \psi \|_{L^4(\tilde{\Omega})}. \quad (2.10)$$

Here the constant $C > 0$ depends only on a , m and $\tilde{\Omega}$.

Proof. The estimate (2.6) is rather standard and is obtained in [16]. It can be derived using a simple energy argument as in [5], without relying on regularity properties of $(\psi, \mathbf{A})$.

Inserting $\phi = \psi$ in (2.2) we get,

$$\int_{\tilde{\Omega}} (w_m(x) |(\nabla - i\kappa H \mathbf{A}) \psi|^2 + \kappa^2 V_a(x) |\psi|^2 + \kappa^2 \mathbf{1}_\Omega(x) |\psi|^4) dx = 0. \quad (2.11)$$

Now (2.7) and (2.8) are consequences of (2.11). The estimate (2.9) is a consequence of (2.6).

Now, we prove (2.10). Up to a gauge transformation, we may assume that $\mathbf{A} \in H_{\text{div}}^1(\psi, \mathbf{A})$. Inserting $\mathbf{a} = \mathbf{A} - \mathbf{F}$ in (2.3) and estimating we get,

$$\begin{aligned} \kappa H \int_{\tilde{\Omega}} |\text{curl}(\mathbf{A} - \mathbf{F})|^2 dx \\ \leq \max \left(1, \frac{1}{m} \right) \|\psi\|_{L^\infty(\tilde{\Omega})} \|(\nabla - i\kappa H \mathbf{A})\psi\|_{L^2(\tilde{\Omega})} \|(\mathbf{A} - \mathbf{F})\psi\|_{L^2(\tilde{\Omega})}. \end{aligned}$$

Invoking (2.6), (2.7) and applying a further Cauchy-Schwarz inequality, we get,

$$\begin{aligned} \kappa H \int_{\tilde{\Omega}} |\text{curl}(\mathbf{A} - \mathbf{F})|^2 dx \leq \\ \kappa \max \left(1, \frac{1}{\sqrt{m}} \right) \|\psi\|_{L^2(\Omega)} \|\psi\|_{L^4(\tilde{\Omega})} \|\mathbf{A} - \mathbf{F}\|_{L^4(\tilde{\Omega})}. \end{aligned} \quad (2.12)$$

Now, by the Sobolev embedding theorem and Lemma 2.1, we get

$$\|\mathbf{A} - \mathbf{F}\|_{L^4(\tilde{\Omega})} \leq C_{\text{Sob}} \|\mathbf{A} - \mathbf{F}\|_{H^1(\Omega)} \leq \tilde{C} \|\text{curl}(\mathbf{A} - \mathbf{F})\|_{L^2(\tilde{\Omega})}.$$

Thus, we can divide through by $\|\text{curl} \mathbf{A} - \mathbf{F}\|_{L^2(\tilde{\Omega})}$ in (2.12) to get (2.10). $\square$

The next theorem gives the finiteness of the critical fields H_{C_3} and is well-known (see [17, 7] for superconductors in vacuum and [16] for a setting similar to ours). We give an easy (spectral) proof for completeness.

Theorem 2.3. *There exists a constant $C > 0$ such that if*

$$\kappa \geq 1 \quad \text{and} \quad H \geq C\kappa, \quad (2.13)$$

then the only weak solution to (2.1) is the normal state $(0, \mathbf{F})$.

Proof. We may assume—after possibly performing a gauge transformation—that the stationary point satisfies $(\psi, \mathbf{A}) \in H^1(\tilde{\Omega}; \mathbb{C}) \times H_{\text{div}}^1(\tilde{\Omega}; \mathbb{R}^2)$. Suppose that $\psi \neq 0$.

Using $\|\psi\|_\infty \leq 1$, we have the pointwise inequality

$$|(\nabla - i\kappa H \mathbf{F})\psi|^2 \leq 2|(\nabla - i\kappa H \mathbf{A})\psi|^2 + 2(\kappa H)^2 |\mathbf{A} - \mathbf{F}|^2. \quad (2.14)$$

Upon integration of (2.14) and using Lemma 2.2, the Sobolev inclusion $H^1 \hookrightarrow L^4$ and Lemma 2.1, we find

$$\int_{\tilde{\Omega}} |(\nabla - i\kappa H \mathbf{F})\psi|^2 dx \leq C\kappa^2 \|\psi\|_{L^2(\tilde{\Omega})}^2, \quad (2.15)$$

for some constant $C > 0$. This implies—since $\psi \neq 0$ by assumption—that the lowest Neumann eigenvalue $\mu^N(\kappa H)$ of $-(\nabla - i\kappa H \mathbf{F})^2$ in $L^2(\tilde{\Omega})$ satisfies

$$\mu^N(\kappa H) \leq C\kappa^2. \quad (2.16)$$

However, since $\tilde{\Omega}$ is smooth, we have (see [7])

$$\mu^N(\kappa H) \geq C'\kappa H, \quad \forall H \geq H_0, \quad (2.17)$$

for positive constants C' and H_0 independent from κ . Combining (2.17) and (2.16) yields the result. $\square$

In the next theorem, we show that the L^4 analogue of (2.8) holds for critical points.

Theorem 2.4. *Let $(\psi, \mathbf{A}) \in H^1(\tilde{\Omega}; \mathbb{C}) \times H_{\text{div}}^1(\tilde{\Omega}; \mathbb{R}^2)$ be a weak solution of (2.1). Then the following estimate holds,*

$$a \int_{\tilde{\Omega} \setminus \Omega} |\psi|^4 dx \leq \int_{\Omega} |\psi|^4 dx.$$

The estimate in Theorem 2.4 is an essential ingredient in proving Theorem 2.7 below (which links the existence of non-trivial critical points to a spectral condition). In order to prove Theorem 2.4, we need to establish some regularity properties for solutions of (2.1).

Lemma 2.5. *Let $(\psi, \mathbf{A}) \in H^1(\tilde{\Omega}; \mathbb{C}) \times H_{\text{div}}^1(\tilde{\Omega}; \mathbb{R}^2)$ be a weak solution of (2.1). Then,*

- (1) $(\psi, \mathbf{A}) \in H^2(\tilde{\Omega} \setminus \partial\Omega; \mathbb{C}) \times H^2(\tilde{\Omega}; \mathbb{R}^2)$.
- (2) $(\psi, \mathbf{A}) \in C^\infty(\tilde{\Omega} \setminus \partial\Omega; \mathbb{C}) \times C^\infty(\tilde{\Omega} \setminus \partial\Omega; \mathbb{R}^2)$.

Let us emphasize that the complete regularity of $(\psi, \mathbf{A})$ up to the boundary of Ω does not follow from standard regularity theory for elliptic PDE, and hence deserves to be studied independently.

Proof of Lemma 2.5. The interior regularity of $(\psi, \mathbf{A})$ (statement (2) above) is obtained through a standard bootstrapping argument, see [29, Proposition 3.8].

We move now to the H^2 regularity. Since $\mathbf{A} \in H_{\text{div}}^1(\tilde{\Omega})$, the equation for $\mathbf{A}$ in (2.1) becomes,

$$-\Delta \mathbf{A} = g(x) \quad \text{in } \tilde{\Omega},$$

coupled with the boundary conditions,

$$\text{curl } \mathbf{A} = 1, \quad \nu \cdot \mathbf{A} = 0, \quad \text{on } \partial\tilde{\Omega}.$$

Here

$$g(x) := (\kappa H)^{-1} w_m(x) (i\psi, (\nabla - i\kappa H \mathbf{A})\psi) \in L^2(\tilde{\Omega}).$$

Now $\mathbf{A} \in H^2(\tilde{\Omega})$ follows from the $W^{k,p}$ -regularity of the curl-div system (see [1]).

To obtain the regularity of ψ one has to be more careful, since ψ satisfies formally a transmission condition on $\partial\Omega$,

$$\mathcal{T}_{\partial\Omega}^{\text{int}}(\nu \cdot (\nabla - i\kappa H \mathbf{A})\psi) = \frac{1}{m} \mathcal{T}_{\partial\Omega}^{\text{ext}}(\nu \cdot (\nabla - i\kappa H \mathbf{A})\psi). \quad (2.18)$$

Here

$$\mathcal{T}_{\partial\Omega}^{\text{int}} : H^1(\Omega) \rightarrow L^2(\partial\Omega), \quad \mathcal{T}_{\partial\Omega}^{\text{ext}} : H^1(\mathbb{R}^2 \setminus \Omega) \rightarrow L^2(\partial\Omega) \quad (2.19)$$

are respectively, the ‘interior’ and ‘exterior’ trace operators, and ν is the outward unit normal vector of $\partial\Omega$.

The crucial point is now to apply a suitable gauge transformation which makes the transmission condition (2.18) independent of the vector potential $\mathbf{A}$.

Let χ be the solution of the following boundary value problem,

$$-\Delta \chi = 0 \quad \text{in } \Omega, \quad \nu \cdot \nabla \chi = \nu \cdot \mathbf{A} \quad \text{on } \partial\Omega.$$

Since $\mathbf{A} \in H^2(\tilde{\Omega})$, standard regularity theory gives $\chi \in W^{2,p}(\Omega)$ for every $p \geq 1$. Let us fix a choice of $p > 2$ such that Sobolev embedding gives $W^{2,p} \subset C^{1,\alpha}$ for some $\alpha \in]0, 1[$.

Given a smooth domain K such that $\bar{\Omega} \subset K \subset \bar{K} \subset \tilde{\Omega}$, we can extend χ to a function $\tilde{\chi} \in W^{2,p}(\tilde{\Omega})$ such that $\text{supp } \chi \subset K$ (see Remark on p.257 in [6]).

Now, defining,

$$\varphi = \psi e^{i\tilde{\chi}}, \quad \mathbf{B} = \mathbf{A} - \nabla \tilde{\chi},$$

the condition (2.18) reads formally,

$$\mathcal{T}_{\partial\Omega}^{\text{int}}(\nu \cdot \nabla \varphi) = \frac{1}{m} \mathcal{T}_{\partial\Omega}^{\text{ext}}(\nu \cdot \nabla \varphi). \quad (2.20)$$

Actually, the equation for φ becomes,

$$-\text{div}(w_m(x) \nabla \varphi) = f \quad \text{in } \tilde{\Omega},$$

with

$$f(x) = -(\kappa H)w_m(x) [2i(\mathbf{B} \cdot \nabla) + i(\operatorname{div} \mathbf{B}) + \kappa H|\mathbf{B}|^2] \varphi - \kappa^2 (V_a(x)\varphi + \mathbf{1}_\Omega(x)|\varphi|^2\varphi).$$

Here w_m and V_a are as in (1.15). Moreover, the equation for φ is supplemented with the boundary condition,

$$\nu \cdot \nabla \varphi = 0 \quad \text{on } \partial\tilde{\Omega}.$$

The obtained equation for φ is of the form studied in [22, Appendix B], where L^2 -type estimates are shown to hold for the solutions. Let us see how we will implement the aforementioned point.

Lemma 2.2 gives $|\varphi| \leq 1$. Moreover, since $\varphi \in H^1(\tilde{\Omega})$, $\mathbf{A} \in H^2(\tilde{\Omega})$ and $\mathbf{B} = \mathbf{A} + \nabla \tilde{\chi}$ is bounded, we deduce that $f \in L^2(\tilde{\Omega})$. Applying now Theorem B.1 in [22], we deduce that $\varphi \in H^2(\tilde{\Omega} \setminus \partial\Omega)$. Since $\psi = \varphi e^{-i\tilde{\chi}}$ and $\tilde{\chi} \in W^{2,p}(\tilde{\Omega}) \subset C^{1,\alpha}(\tilde{\Omega})$, then $\psi \in H^2(\tilde{\Omega} \setminus \partial\Omega)$. $\square$

Proof of Theorem 2.4. Thanks to Lemma 2.5, $(\psi, \mathbf{A}) \in H^1(\tilde{\Omega}; \mathbb{C}) \times H_{\operatorname{div}}^1(\tilde{\Omega}; \mathbb{R}^2)$ being a solution of (2.1), the function $\psi \in C^\infty(\tilde{\Omega} \setminus \partial\Omega)$, and $(\psi, \mathbf{A})$ is a strong solution of (2.1) in $\tilde{\Omega} \setminus \partial\Omega$.

Consequently, the function,

$$u = |\psi|^2 = \psi \bar{\psi} \in C^\infty(\tilde{\Omega} \setminus \partial\Omega; \mathbb{R}).$$

It is easy to verify that,

$$\frac{1}{2} \Delta |\psi|^2 = \operatorname{Re}(\bar{\psi}(\nabla - i\kappa H \mathbf{A})^2 \psi) + |(\nabla - i\kappa H \mathbf{A})\psi|^2, \quad \forall x \in \tilde{\Omega} \setminus \partial\Omega.$$

Thus, defining the function,

$$f(x) = \frac{1}{2} w_m(x) |(\nabla - i\kappa H \mathbf{A})\psi(x)|^2,$$

we get that $u = |\psi|^2$ is a strong solution in $\tilde{\Omega} \setminus \partial\Omega$ of the equation,

$$-\frac{1}{2} \operatorname{div}(w_m \nabla u) + \kappa^2 (V_a + \mathbf{1}_\Omega u) u + f = 0. \quad (2.21)$$

Here we remind the reader that the functions w_m and V_a are defined in (1.15).

Let us show that u is a weak solution of (2.21) in $\tilde{\Omega}$. To that end, we need only verify that

$$\mathcal{T}_{\partial\Omega}^{\operatorname{int}}(\nu \cdot \nabla u) = \frac{1}{m} \mathcal{T}_{\partial\Omega}^{\operatorname{ext}}(\nu \cdot \nabla u) \quad \text{in } L^2(\partial\Omega), \quad (2.22)$$

where $\mathcal{T}_{\partial\Omega}^{\operatorname{int}}$ and $\mathcal{T}_{\partial\Omega}^{\operatorname{ext}}$ are the trace operators introduced in (2.19), and ν is the outward unit normal vector of $\partial\Omega$.

Notice that $\nabla u = 2 \operatorname{Re}(\psi \nabla \bar{\psi}) \in H^1(\tilde{\Omega} \setminus \partial\Omega)$ since $\psi \in H^2(\tilde{\Omega} \setminus \partial\Omega; \mathbb{C})$ and $\tilde{\Omega} \subset \mathbb{R}^2$, hence the trace of $\nu \cdot \nabla u$ is well defined in the usual sense.

Now, it is easy to verify that,

$$\nabla u = 2 \operatorname{Re}(\psi \nabla \bar{\psi}) = 2 \operatorname{Re}(\psi \overline{(\nabla - i\kappa H \mathbf{A})\psi}).$$

On the other hand, since $(\psi, \mathbf{A}) \in H^2(\tilde{\Omega} \setminus \partial\Omega; \mathbb{C}) \times H^2(\tilde{\Omega}; \mathbb{R}^2)$ and weak solutions of (2.1), they satisfy in particular,

$$\mathcal{T}_{\partial\Omega}^{\operatorname{int}}(\nu \cdot (\nabla - i\kappa H \mathbf{A})\psi) = \frac{1}{m} \mathcal{T}_{\partial\Omega}^{\operatorname{ext}}(\nu \cdot (\nabla - i\kappa H \mathbf{A})\psi)$$

in $L^2(\partial\Omega)$. This shows that (2.22) also holds in $L^2(\partial\Omega)$ and consequently $u = |\psi|^2$ is a weak solution in $\tilde{\Omega}$ of (2.21).

Therefore, multiplying (2.21) by u and integrating, we get,

$$\int_{\tilde{\Omega}} \left(\frac{1}{2} w_m(x) |\nabla u|^2 + V_a(x) u^2 + 1_{\Omega}(x) u^3 + f(x) u \right) dx = 0,$$

with $u \geq 0$ and $f \geq 0$. This yields the estimate of Lemma 2.4. $\square$

2.2. Weak Decay Estimate and applications. Just as in [3], we can derive the following weak decay estimate.

Lemma 2.6. *Assume that $a > 0$ and $m > 0$. There exist positive constants C and C' such that, if $(\psi, \mathbf{A})$ is a solution of (2.1) with:*

$$\kappa(H - \kappa) \geq C,$$

then the following estimate holds:

$$\begin{aligned} \max \left(1, \frac{a\kappa}{H - \kappa} \right) \|\psi\|_{L^2(\tilde{\Omega})}^2 &\leq C \int_{\{x \in \tilde{\Omega}: \sqrt{\kappa(H - \kappa)} \operatorname{dist}(x, \partial\Omega) \leq 1\}} |\psi(x)|^2 dx \\ &\leq \frac{C'}{\kappa(H - \kappa)}. \end{aligned}$$

Proof. The last inequality is an easy consequence of (2.6), so we will only establish the first one.

Let $\chi \in C^\infty(\mathbb{R})$ be a standard cut-off function such that,

$$\chi = 1 \quad \text{in } [1, \infty[, \quad \chi = 0 \quad \text{in }]-\infty, 1/2].$$

Define $\lambda = 1/\sqrt{\kappa(\kappa - H)}$, and

$$\chi_\lambda(x) = \chi \left(\frac{\operatorname{dist}(x, \partial\Omega)}{\lambda} \right), \quad x \in \tilde{\Omega}.$$

A simple calculation yields the following localization formula:

$$\begin{aligned} \int_{\tilde{\Omega}} w_m(x) (|(\nabla - i\kappa H \mathbf{A}) \chi_\lambda \psi|^2 - |\nabla \chi_\lambda|^2 |\psi|^2) dx \\ = \operatorname{Re} \int_{\tilde{\Omega}} \left(w_m(x) \overline{(\nabla - i\kappa H \mathbf{A}) \psi} \cdot (\nabla - i\kappa H \mathbf{A}) (\chi_\lambda^2 \psi) \right) dx, \end{aligned}$$

where w_m is introduced in (1.15).

Now, we use the weak formulation of the first G-L equation in (2.1) and get,

$$\begin{aligned} \int_{\tilde{\Omega}} w_m(x) (|(\nabla - i\kappa H \mathbf{A}) \chi_\lambda \psi|^2 - |\nabla \chi_\lambda|^2 |\psi|^2) dx + a\kappa^2 \int_{\tilde{\Omega} \setminus \Omega} \chi_\lambda^2 |\psi|^2 dx \\ = \kappa^2 \int_{\Omega} \chi_\lambda^2 (1 - |\psi|^2) |\psi|^2 dx. \end{aligned} \quad (2.23)$$

The next step is to give a lower bound to the first term on the left hand side of (2.23). This is done via an elementary inequality from the spectral theory of magnetic Schrödinger operators (see e.g. [2, Thm 2.9] or [7, Lemma 2.4.1]). Actually, since the function χ_λ has compact support not meeting the boundary $\partial\Omega$, and $w_m = 1$ in Ω , the following inequality holds,

$$\int_{\tilde{\Omega}} w_m(x) |(\nabla - i\kappa H \mathbf{A}) \chi_\lambda \psi|^2 dx \geq \kappa H \int_{\Omega} (\operatorname{curl} \mathbf{A}) |\chi_\lambda \psi|^2 dx.$$

Writing $\operatorname{curl} \mathbf{A} = 1 + (\operatorname{curl} \mathbf{A} - 1)$ then applying a Cauchy-Schwarz inequality, we get,

$$\begin{aligned} \int_{\tilde{\Omega}} w_m(x) |(\nabla - i\kappa H \mathbf{A}) \chi_\lambda \psi|^2 dx &\geq \kappa H \int_{\Omega} w_m(x) |\chi_\lambda \psi|^2 dx \\ &\quad - \kappa H \|\operatorname{curl} \mathbf{A} - 1\|_{L^2(\Omega)} \|\chi_\lambda \psi\|_{L^4(\Omega)}^2. \end{aligned}$$

Implementing the estimates (2.10), (2.9) and (2.8) we get, for some constant $c > 0$, the following lower bound,

$$\begin{aligned} \int_{\tilde{\Omega}} w_m(x) |(\nabla - i\kappa H \mathbf{A}) \chi_\lambda \psi|^2 dx \\ \geq \kappa H \int_{\tilde{\Omega}} |\chi_\lambda \psi|^2 dx - c\kappa \|\psi\|_{L^2(\Omega)} \|\chi_\lambda \psi\|_{L^4(\Omega)}^2. \end{aligned} \quad (2.24)$$

Upon substitution in (2.23) and a rearrangement of terms, we deduce,

$$\begin{aligned} \kappa(H - \kappa) \int_{\tilde{\Omega}} |\chi_\lambda \psi|^2 dx + a\kappa^2 \int_{\tilde{\Omega} \setminus \Omega} |\chi_\lambda \psi|^2 dx \leq c\kappa \|\psi\|_{L^2(\Omega)} \|\chi_\lambda \psi\|_{L^4(\Omega)}^2 \\ + \|\chi'\|_{L^\infty(\mathbb{R})}^2 \lambda^{-2} \int_{\{\text{dist}(x, \partial\Omega) \leq \lambda\}} w_m(x) |\psi|^2 dx - \kappa^2 \int_{\tilde{\Omega}} \chi_\lambda^2 |\psi|^4 dx. \end{aligned}$$

Implementing again a Cauchy-Schwarz inequality, we get

$$\begin{aligned} \kappa(H - \kappa) \int_{\tilde{\Omega}} |\chi_\lambda \psi|^2 dx + a\kappa^2 \int_{\tilde{\Omega} \setminus \Omega} |\chi_\lambda \psi|^2 dx \\ \leq c^2 \|\psi\|_{L^2(\Omega)}^2 + \|\chi'\|_{L^\infty(\mathbb{R})}^2 \lambda^{-2} \int_{\{\text{dist}(x, \partial\Omega) \leq \lambda\}} w_m(x) |\psi|^2 dx \\ + \kappa^2 \int_{\tilde{\Omega}} (\chi_\lambda^4 - \chi_\lambda^2) |\psi|^4 dx, \end{aligned}$$

where the last term on the right hand side above is negative, since $0 \leq \chi_\lambda \leq 1$.

Decomposing the integral $\int_{\tilde{\Omega}} |\psi|^2 = \int_{\tilde{\Omega}} |\chi_\lambda \psi|^2 + \int_{\tilde{\Omega}} (1 - \chi_\lambda^2) |\psi|^2$, and assuming that

$$\kappa(H - \kappa) \geq 2c^2,$$

we get

$$\begin{aligned} \frac{1}{2} \max(\kappa(H - \kappa), a\kappa^2) \int_{\tilde{\Omega}} |\chi_\lambda \psi|^2 dx \\ \leq \left(c^2 + \max\left(1, \frac{1}{m}\right) \|\chi'\|_{L^\infty(\mathbb{R})}^2 \lambda^{-2} \right) \int_{\{\text{dist}(x, \partial\Omega) \leq \lambda\}} |\psi|^2 dx. \end{aligned}$$

Recall that $\lambda = 1/\sqrt{\kappa(H - \kappa)}$. The conditions on χ and $\kappa(H - \kappa)$ imply that

$$\max\left(1, \frac{a\kappa}{H - \kappa}\right) \int_{\tilde{\Omega}} |\chi_\lambda \psi|^2 dx \leq 4 \max\left(1, \frac{1}{m}\right) \|\chi'\|_{L^\infty(\mathbb{R})}^2 \int_{\{\text{dist}(x, \partial\Omega) \leq \lambda\}} |\psi|^2 dx.$$

Consequently, we get,

$$\begin{aligned} \max\left(1, \frac{a\kappa}{H - \kappa}\right) \int_{\tilde{\Omega}} |\psi|^2 dx \\ \leq \left(4 \max\left(1, \frac{1}{m}\right) \|\chi'\|_{L^\infty(\mathbb{R})}^2 + 1 \right) \int_{\{\text{dist}(x, \partial\Omega) \leq \lambda\}} |\psi|^2 dx. \end{aligned}$$

Choosing $C = \max\left(c^2, 4 \max\left(1, \frac{1}{m}\right) \|\chi'\|_{L^\infty(\mathbb{R})}^2 + 1\right)$, we get the desired bound. $\square$

The next theorem gives a purely spectral criterion for the existence of non-trivial critical points of (1.1).

Theorem 2.7. *Given $a > 0$ and $m > 1$, there exists $\kappa_0 > 0$ such that, for all $\kappa \geq \kappa_0$, the following two statements are equivalent:*

- (1) *There exists a solution $(\psi, \mathbf{A})$ of (2.1) with $\|\psi\|_{L^2(\tilde{\Omega})} \neq 0$.*
- (2) *The parameters κ and H satisfy $\mu^{(1)}(\kappa, H) < 0$, where the eigenvalue $\mu^{(1)}(\kappa, H)$ is introduced in (1.7).*

Proof. It is well known that the second statement implies the first one. Actually, we only use $(t\psi_*, 0)$, with t sufficiently small and ψ_* an eigenfunction associated with $\mu^{(1)}(\kappa, H)$, as a test configuration for the functional (1.1). The resulting energy will be lower than that of a normal state. Hence a minimizer, which is a solution of (2.1), will be non-trivial.

We assume now that the first statement holds and we show that the second statement is true provided that κ is sufficiently large.

Thanks to Theorem 2.3, it is sufficient to deal with applied magnetic fields satisfying $H \leq C\kappa$, for some constant $C > 0$.

On the other hand, if $c > 0$ is a sufficiently small constant, then we can show that

$$\mu^{(1)}(\kappa, H) < 0 \quad \text{for all } H < c\kappa. \quad (2.25)$$

In order to see this, notice that the variational min-max principle gives,

$$\mu^{(1)}(\kappa, H) \leq \mu^D(\kappa H; \Omega) - \kappa^2,$$

where $\mu^D(\kappa H; \Omega)$ is the lowest Dirichlet eigenvalue of $-(\nabla - i\kappa H \mathbf{F})^2$ in $L^2(\Omega)$. Standard estimates on Dirichlet realizations of magnetic operators (see [18]) yield the existence of a constant $\tilde{C} > 0$ such that

$$\mu^D(B; \Omega) \leq \tilde{C} \max(B, 1) \quad \forall B > 0,$$

implying (2.25).

Therefore, we restrict ourselves to applied magnetic fields satisfying,

$$c\kappa \leq H \leq C\kappa.$$

Using Lemma 2.6 and a Cauchy-Schwarz inequality, we find a positive constant $C > 0$ such that

$$\|\psi\|_{L^2(\tilde{\Omega})}^2 \leq C \left(\int_{\{\text{dist}(x, \partial\Omega) \leq \frac{1}{\kappa}\}} dx \right)^{1/2} \|\psi\|_{L^4(\tilde{\Omega})}^2 \leq \frac{C'}{\sqrt{\kappa}} \|\psi\|_{L^4(\Omega)}^2, \quad (2.26)$$

where we use Theorem 2.4 to get the last inequality.

Since $(\psi, \mathbf{A})$ is a (weak) solution of (2.1), we get by setting $\phi = \psi$ in (2.2) together with the assumption on ψ that

$$0 < \kappa^2 \|\psi\|_{L^4(\Omega)}^4 \leq - \int_{\tilde{\Omega}} (w_m(x) |(\nabla - i\kappa H \mathbf{A})\psi|^2 + \kappa^2 V_a(x) |\psi|^2) dx =: \Delta, \quad (2.27)$$

where w_m and V_a are introduced in (1.15). Notice that use (2.8) implies that $\|\psi\|_{L^4(\Omega)} \neq 0$.

Implementing (2.27) in (2.26), we get

$$\|\psi\|_{L^2(\tilde{\Omega})}^2 \leq C'' \sqrt{\Delta} \kappa^{-3/2}. \quad (2.28)$$

We estimate

$$\begin{aligned} & |(\nabla - i\kappa H \mathbf{A})\psi|^2 \\ & \geq (1 - \sqrt{\Delta} \kappa^{-3/4}) |(\nabla - i\kappa H \mathbf{F})\psi|^2 - \frac{1}{\sqrt{\Delta} \kappa^{-3/4}} (\kappa H)^2 |(\mathbf{A} - \mathbf{F})\psi|^2. \end{aligned}$$

This yields the following estimate on Δ :

$$\begin{aligned} \Delta & \leq \left(-\mu^{(1)}(\kappa, H) + \min \left(1, \frac{1}{m} \right) \sqrt{\Delta} \kappa^{-3/4} \mu^N(\kappa H) \right) \|\psi\|_{L^2(\tilde{\Omega})}^2 \\ & \quad + \min \left(1, \frac{1}{m} \right) \frac{\kappa^{3/4}}{\sqrt{\Delta}} (\kappa H)^2 \int_{\tilde{\Omega}} |(\mathbf{A} - \mathbf{F})\psi|^2 dx, \end{aligned} \quad (2.29)$$

where $\mu^N(\kappa H)$ is the lowest eigenvalue of the magnetic Neumann Laplacian $-(\nabla - i\kappa H \mathbf{F})^2$ in $\tilde{\Omega}$.

We recall that $H = \mathcal{O}(\kappa)$. Using the asymptotic behavior of $\mu^N(B)$ as $B \rightarrow \infty$ together with the estimate (2.28), we get:

$$\sqrt{\Delta} \kappa^{-3/4} \mu^N(\kappa H) \|\psi\|_{L^2(\tilde{\Omega})}^2 \leq C \kappa^{-1/4} \Delta. \quad (2.30)$$

On the other hand, using a Cauchy-Schwarz inequality and the estimate (2.27), we get:

$$(\kappa H)^2 \int_{\tilde{\Omega}} |(\mathbf{A} - \mathbf{F})\psi|^2 dx \leq (\kappa H)^2 \|\mathbf{A} - \mathbf{F}\|_{L^4(\tilde{\Omega})}^2 \frac{\sqrt{2\Delta}}{\kappa}.$$

By Sobolev embedding and Lemma 2.1,

$$(\kappa H)^2 \|\mathbf{A} - \mathbf{F}\|_{L^4(\tilde{\Omega})}^2 \leq C(\kappa H)^2 \|\operatorname{curl} \mathbf{A} - 1\|_{L^2(\tilde{\Omega})}^2 \leq C \kappa^2 \|\psi\|_{L^4(\tilde{\Omega})}^4.$$

Using again Theorem 2.4, we deduce that,

$$(\kappa H)^2 \|\mathbf{A} - \mathbf{F}\|_{L^4(\tilde{\Omega})}^2 \leq C \kappa^2 \|\psi\|_{L^4(\Omega)}^4 \leq C \Delta.$$

Therefore, implementing all the above estimates in (2.29), we get

$$\Delta \leq -\mu^{(1)}(\kappa, H) \|\psi\|_{L^2(\tilde{\Omega})}^2 + C \frac{\Delta}{\kappa^{1/4}}.$$

Knowing that $\Delta > 0$ (see (2.27)), we deduce for κ sufficiently large the desired inequality, $\mu^{(1)}(\kappa, H) < 0$. $\square$

Proof of Theorem 1.1. We can now finish the proof of Theorem 1.1 as follows. Theorem 2.7 gives $\mathcal{N}^{\text{sc}}(a, m; \kappa) = \mathcal{N}^{\text{loc}}(a, m; \kappa)$ for κ sufficiently large. On the other hand, we have the trivial inclusions $\mathcal{N}(a, m; \kappa) \subset \mathcal{N}^{\text{sc}}(a, m; \kappa)$ and $\mathcal{N}^{\text{loc}}(a, m; \kappa) \subset \mathcal{N}(a, m; \kappa)$. $\square$

3. MONOTONICITY OF THE FIRST EIGENVALUE

3.1. The constant $\alpha_0(a, m)$. We recall in this section the definition of the constant $\alpha_0(a, m)$ introduced in [23], together with the main properties of a family of ordinary differential operators.

Consider the space $B^k(\mathbb{R}) = H^k(\mathbb{R}) \cap L^2(\mathbb{R}; |t|^k dt)$, $k \in \mathbb{N}$. Given $a, m, \alpha > 0$ and $\xi \in \mathbb{R}$, let us define the quadratic form :

$$B^1(\mathbb{R}) \ni u \mapsto q[a, m, \alpha; \xi](u), \quad (3.1)$$

where :

$$\begin{aligned} q[a, m, \alpha; \xi](u) &= \int_{\mathbb{R}_+} (|u'(t)|^2 + |(t - \xi)u(t)|^2 - \alpha|u(t)|^2) dt \\ &\quad + \int_{\mathbb{R}_-} \left(\frac{1}{m} [|u'(t)|^2 + |(t - \xi)u(t)|^2] + a\alpha|u(t)|^2 \right) dt. \end{aligned} \quad (3.2)$$

We denote by $H[a, m, \alpha; \xi]$ the self-adjoint operator associated to the closed symmetric quadratic form (3.1). The domain of $H[a, m, \alpha; \xi]$ is defined by :

$$D(H[a, m, \alpha; \xi]) = \left\{ u \in B^1(\mathbb{R}); \quad u|_{\mathbb{R}_{\pm}} \in B^2(\mathbb{R}_{\pm}), \quad u'(0_+) = \frac{1}{m} u'(0_-) \right\}, \quad (3.3)$$

and for $u \in D(H[a, m, \alpha; \xi])$, we have,

$$(H[a, m, \alpha; \xi]u)(t) = \begin{cases} [(-\partial_t^2 + (t - \xi)^2 - \alpha)u](t); & \text{if } t > 0, \\ [(\frac{1}{m} \{-\partial_t^2 + (t - \xi)^2\} + a\alpha)u](t); & \text{if } t < 0. \end{cases} \quad (3.4)$$

We denote by $\mu_1(a, m, \alpha; \xi)$ the first eigenvalue of $H[a, m, \alpha; \xi]$ which is given by the min-max principle,

$$\mu_1(a, m, \alpha; \xi) = \inf_{u \in B^1(\mathbb{R}), u \neq 0} \frac{q[a, m, \alpha; \xi](u)}{\|u\|_{L^2(\mathbb{R})}^2}. \quad (3.5)$$

We summarize in the next theorem the main results obtained in [23] concerning the above family of operators.

Theorem 3.1. *The following assertions hold.*

- (1) *Given $a > 0$ and $m > 0$, there exists a unique $\alpha_0(a, m) > 0$ such that*

$$\inf_{\xi \in \mathbb{R}} \mu_1(a, m, \alpha_0(a, m); \xi) = 0.$$

Moreover, $\alpha_0(a, m) = 1$ if $m \leq 1$, and there exists a universal constant $\Theta_0 \in]0, 1[$ such that $\Theta_0 < \alpha_0(a, m) < 1$ if $m > 1$, and

$$\lim_{m \rightarrow \infty} \alpha_0(a, m) = \Theta_0.$$

- (2) *If $0 < \alpha < \alpha_0(a, m)$, then $\inf_{\xi \in \mathbb{R}} \mu_1(a, m, \alpha; \xi) > 0$.*
 (3) *Given $a > 0$, there exists $m_0 > 1$ such that for all $m \geq m_0$ the function $\mathbb{R} \ni \xi \mapsto \mu_1(a, m, \alpha_0(a, m); \xi)$ admits a unique non-degenerate minimum.*
 (4) *For all $a > 0$ and $m \geq m_0$, there exists $\epsilon_0(m) > 0$ such that if*

$$\alpha \in [\alpha_0(a, m) - \epsilon_0(m), \alpha_0(a, m) + \epsilon_0(m)],$$

then the function $\mathbb{R} \ni \xi \mapsto \mu_1(a, m, \alpha; \xi)$ admits a unique non-degenerate minimum, denoted by $\xi(a, m, \alpha)$.

- (5) *For all $m > 0$ and $a > 0$, $\inf_{\xi \in \mathbb{R}} \mu_1(a, m, \alpha; \xi) \rightarrow 0$ as $\alpha \rightarrow \alpha_0(a, m)$.*

Let us mention that the universal constant Θ_0 is the infimum of the spectrum of the Neumann Schrödinger operator with unit magnetic field in $\mathbb{R} \times \mathbb{R}_+$, and it holds that $\frac{1}{2} < \Theta_0 < 1$. We point also that the existence of the constant $m_0 > 1$ in Theorem 3.1 is nontrivial, and is due to a fine asymptotic analysis of the eigenvalue (3.5) as $m \rightarrow \infty$, see [23, Section 3.4].

For further use, we introduce the constant

$$\beta(a, m, \alpha) = \inf_{\xi \in \mathbb{R}} \mu_1(a, m, \alpha; \xi), \quad (3.6)$$

and the set

$$M(a, m, \alpha) = \{\xi \in \mathbb{R} : \mu_1(a, m, \alpha; \xi) = \beta(a, m, \alpha)\}. \quad (3.7)$$

We notice as a result of Theorem 3.1 that $\beta(a, m, \alpha) = 0$ if and only if $\alpha = \alpha_0(a, m)$, and in this case $M(a, m, \alpha) = \{\xi(a, m, \alpha)\}$ when $m \geq m_0$.

Let us also notice that it results from a simple application of the min-max principle together with well known results concerning the harmonic oscillator in $\mathbb{R}_+$,

$$\beta(a, m, \alpha) + \alpha \geq \min \left(\Theta_0, \frac{\Theta_0}{m} + (a+1)\alpha \right), \quad \forall \alpha > 0. \quad (3.8)$$

Furthermore, let $f_{\alpha, \xi}^{a, m}$ be the normalized eigenfunction associated with the eigenvalue (3.5). We introduce the constants

$$C_1(a, m, \alpha; \xi) = \int_{\mathbb{R}_+} (t - \xi)^3 |f_{\alpha, \xi}^{a, m}(t)|^2 dt + \frac{1}{m} \int_{\mathbb{R}_-} (t - \xi)^3 |f_{\alpha, \xi}^{a, m}(t)|^2 dt \quad (3.9)$$

$$- \frac{1}{2} \left(1 - \frac{1}{m} \right) |f_{\alpha, \xi}^{a, m}(0)|^2, \quad (3.10)$$

$$b_1(a, m, \alpha; \xi) = \int_{\mathbb{R}_+} |f_{\alpha, \xi}^{a, m}(t)|^2 dt - a \int_{\mathbb{R}_-} |f_{\alpha, \xi}^{a, m}(t)|^2 dt. \quad (3.11)$$

If $m \geq m_0$, α fills the hypotheses of assertion (4) in Theorem 3.1 and $\xi = \xi(a, m, \alpha)$, then we write simply:

$$C_1(a, m, \alpha) = C_1(a, m, \alpha; \xi(a, m, \alpha)). \quad (3.12)$$

$$b_1(a, m, \alpha) = b_1(a, m, \alpha; \xi(a, m, \alpha)), \quad (3.13)$$

and if further, $\alpha = \alpha_0 = \alpha_0(a, m)$, we introduce the constant (appearing in the asymptotic formula (1.13))

$$C_1(a, m) = -\frac{C_1(a, m, \alpha_0)}{b_1(a, m, \alpha_0)}. \quad (3.14)$$

Finally, we point out that it is proved in [23, Section 3],

$$\lim_{m \rightarrow \infty} C_1(a, m, \alpha_0) = -(1 + 6a\Theta_0^2)C_1^*, \quad \lim_{m \rightarrow \infty} b_1(a, m, \alpha_0) = 1,$$

where $C_1^* > 0$ is a universal constant (see [23, Remark 1.4]). Thus, for large values of m , the constant in (3.12) is negative.

Now, for $m > 1$ and $\alpha > 0$, we call $(\mathcal{P}_{m, \alpha})$ the property below,

$$(\mathcal{P}_{m, \alpha}) \quad \begin{cases} M(a, m, \alpha) = \{\xi(a, m, \alpha)\} \text{ and } \xi(a, m, \alpha) > 0, \\ \xi(a, m, \alpha) \text{ is a non-degenerate minimum point,} \\ C_1(a, m, \alpha) < 0 \quad \text{and} \quad b_1(a, m, \alpha) > 0. \end{cases}$$

We introduce the constant $m_* \geq 1$,

$$m_* = \inf\{m > 1 : \forall m' \geq m, (\mathcal{P}_{m', \alpha}) \text{ holds for } \alpha = \alpha(a, m')\}. \quad (3.15)$$

In view of the result of Theorem 3.1, we may define m_* as above. We emphasize also that the constant m_* depends on a , but we omit that from the notation for the sake of simplicity. It is conjectured in [23] that $m_* = 1$ for all $a > 0$.

Now, given $a > 0$ and $m > m_*$, we introduce

$$\epsilon_*(m) = \sup\{\epsilon \in]0, d_0/2[: \forall \alpha \in [\alpha_0(a, m) - \epsilon, \alpha_0(a, m) + \epsilon], (\mathcal{P}_{m, \alpha}) \text{ holds}\}, \quad (3.16)$$

where $d_0 := \Theta_0 - \frac{1}{2} > 0$.

For the special case of a disc domain, one more constant will appear to be relevant (this is $C_2(a, m, \alpha)$ introduced below). Given $a > 0$ and $m > m_*$, the lowest eigenvalue of the operator (3.4) for $\xi = \xi(a, m, \alpha)$ is $\beta(a, m, \alpha)$ introduced in (3.6). Hence, the regularized resolvent, i.e. the operator $(\xi = \xi(a, m, \alpha))$,

$$R_0[a, m, \alpha](\phi) = \begin{cases} [H[a, m, \alpha; \xi] - \beta(a, m, \alpha)]^{-1} \phi, & \text{if } \phi \perp f_{\alpha, \xi}^{a, m}, \\ 0, & \text{otherwise,} \end{cases} \quad (3.17)$$

is bounded in $L^2(\mathbb{R})$. Letting $f = f_{\alpha, \xi}^{a, m}$ for $\xi = \xi(a, m, \alpha)$, $\tilde{w}_m(t) = 1$ if $t > 0$ and $\tilde{w}_m(t) = \frac{1}{m}$ if $t < 0$, then it is proved in [23, Proposition 3.6] that the functions f and $\tilde{w}_m f$ are orthogonal in $L^2(\mathbb{R})$, hence the integral

$$I_2(a, m, \alpha) = \int_0^\infty (t - \xi) f R_0(t - \xi) f dt + \frac{1}{m^2} \int_{-\infty}^0 (t - \xi) f R_0(t - \xi) f dt$$

is positive. Notice that we write simply R_0 for the operator (3.17). Now, we introduce the constant,

$$C_2(a, m, \alpha) = \int_0^\infty |f(t)|^2 dt + \frac{1}{m} \int_{-\infty}^0 |f(t)|^2 dt - 4I_2(a, m, \alpha). \quad (3.18)$$

Since $m > 1$ and $\int_{\mathbb{R}} |f|^2 dt = 1$, it is clear that $C_2(a, m, \alpha) < 1$. Recalling that

$$\alpha_0(a, m) > \Theta_0 > \frac{1}{2},$$

we get for $\alpha \in [\alpha_0(a, m) - \epsilon_*(m), \alpha_0(a, m) + \epsilon_*(m)]$ that

$$\alpha_0(a, m) - \frac{1}{2} C_2(a, m, \alpha) > \frac{d_0}{2}, \quad d_0 := \Theta_0 - \frac{1}{2}. \quad (3.19)$$

3.2. Notation, hypotheses and announcement of Main Result. We assume that $\Omega \subset \mathbb{R}^2$, $\tilde{\Omega} \subset \mathbb{R}^2$ are open, bounded and have smooth boundaries such that

$$\overline{\Omega} \subset \tilde{\Omega}. \quad (3.20)$$

Given $B > 0$ and $\alpha > 0$, we denote by $P[B, \alpha]$ the self-adjoint operator in $L^2(\tilde{\Omega})$ generated by the quadratic form

$$H_{B\mathbf{F}}^1(\tilde{\Omega}) \ni \phi \mapsto Q[B, \alpha](\phi) = \int_{\tilde{\Omega}} (w_m(x)|(\nabla - iB\mathbf{F})\phi|^2 + \alpha BV_a(x)|\phi|^2) dx, \quad (3.21)$$

where w_m and V_a are introduced in (1.15), and

$$H_{B\mathbf{F}}^1(\tilde{\Omega}) = \{u \in L^2(\tilde{\Omega}) : (\nabla - iB\mathbf{F})u \in L^2(\mathbb{R}^2)\}. \quad (3.22)$$

Notice that since $\tilde{\Omega}$ is bounded, $H_{B\mathbf{F}}^1(\tilde{\Omega}) = H^1(\tilde{\Omega})$ and hence the form domain of Q is independent of B .

We introduce further the lowest eigenvalue of the operator $P[B, \alpha]$:

$$\mu_1(B, \alpha) = \inf_{\substack{\phi \in H^1(\tilde{\Omega}) \\ \phi \neq 0}} \frac{Q[B, \alpha](\phi)}{\|\phi\|_{L^2(\tilde{\Omega})}^2}. \quad (3.23)$$

By taking $B = \kappa H$ and $\alpha = \kappa/H$, we see the connection with the critical fields introduced in (1.5):

$$\mu^{(1)}(\kappa, H) = \mu_1(B, \alpha). \quad (3.24)$$

Furthermore, we put,

$$\lambda_1(B, \alpha) = \mu_1(B, \alpha) + \alpha B. \quad (3.25)$$

As an application of standard analytic perturbation theory, we have the following proposition.

Proposition 3.2. *Given $a > 0$, $m > 0$ and $\kappa > 0$, the following one sided derivatives*

$$\begin{aligned} \partial_H \mu_{\pm}^{(1)}(\kappa, H) &= \lim_{\epsilon \rightarrow 0_{\pm}} \frac{\mu^{(1)}(\kappa, H + \epsilon) - \mu^{(1)}(\kappa, H)}{\epsilon}, \\ \partial_B \lambda_{1,\pm}(B, \alpha) &= \lim_{\epsilon \rightarrow 0_{\pm}} \frac{\lambda_1(B + \epsilon, \alpha) - \lambda_1(B, \alpha)}{\epsilon}, \end{aligned}$$

exist for all $H > 0$, $B > 0$ and $\alpha > 0$, and

$$\partial_B \lambda_{1,+}(B, \alpha) \leq \partial_B \lambda_{1,-}(B, \alpha).$$

Moreover, there exist L^2 -normalized ground states φ_{H+} and φ_{H-} associated with $\mu^{(1)}(\kappa, H)$ such that,

$$\partial_H \mu_{\pm}^{(1)}(\kappa, H) = \kappa \left(\partial_B \lambda_{1,\pm}(B, \alpha) \Big|_{\substack{B=\kappa H \\ \alpha=\kappa/H}} - \frac{\kappa}{H} \int_{\tilde{\Omega}} (V_a(x) + 1) |\varphi_{H\pm}(x)|^2 dx \right). \quad (3.26)$$

Here V_a is introduced in (1.15).

Proof. Let us explain briefly why the one sided derivatives $\partial_B \lambda_{1,\pm}(B, \alpha)$ above exist; the one sided derivatives in H exist for exactly the same reason.

As we already mentioned, for real B , the operator $P[B, \alpha]$ is self adjoint, has compact resolvent and its form domain, $H^1(\tilde{\Omega})$, is independent from B . Then, one can show that there exists z_0 sufficiently small such that the operator

$$P[z, \alpha] = \nabla_{z\mathbf{F}} \cdot w_m \nabla_{z\mathbf{F}} + \alpha z V_a$$

is analytic of type (B) in² $D_0 = \{z = x + iy : x > 0, |y| < z_0\}$, see [25, p. 392] for the definition of type (B) operators. In particular, for a given $B > 0$, the eigenvalue $\mu_1(B, \alpha)$ has finite multiplicity n . Now, by analytic perturbation theory (see [25, Theorem 4.2, p. 395]) applied to the family of operators $D_0 \ni z \mapsto P[z, \alpha]$, we get the existence of $\epsilon > 0$ and $2n$ analytic functions

$$(B - \epsilon, B + \epsilon) \ni z \mapsto \phi_j(z) \in H^1(\tilde{\Omega}) \setminus \{0\}, \quad (B - \epsilon, B + \epsilon) \ni z \mapsto E_j(z) \in \mathbb{R}$$

such that

$$P[z, \alpha](\phi_j(z)) = E_j(z)\phi_j(z), \quad E_j(B) = \mu_1(B, \alpha).$$

By choosing $\epsilon > 0$ sufficiently small, we get the existence of $j_{\pm} \in \{1, 2, \dots, n\}$ such that

$$\begin{aligned} E_{j_+}(z) &= \min_{j \in \{1, 2, \dots, n\}} E_j(z) \quad \text{for } B < z < B + \epsilon, \\ E_{j_-}(z) &= \min_{j \in \{1, 2, \dots, n\}} E_j(z) \quad \text{for } B - \epsilon < z < B. \end{aligned}$$

With this choice, it is clear that $\partial_B \lambda_{1,\pm}(B, \alpha) = E'_{j_{\pm}}(B) + \alpha$.

Furthermore, we can choose the eigenfunctions $\phi_{j_{\pm}}(z)$ to be L^2 -normalized. Now, the equality (3.26) is obtained through differentiation of the relation (in $z = 0_{\pm}$)

$$\lambda_1(B + z, \alpha) = Q[B + z, \alpha](\phi_{j_{\pm}}(B + z)) + \alpha(B + z),$$

and the application of the chain rule. $\square$

In this section, we shall work under the following hypothesis on the constant m :

$$m > m_*, \quad (3.27)$$

where $m_* > 1$ is the constant introduced in (3.15).

The next hypothesis is on the constant α :

$$-\epsilon_*(m) \leq \alpha - \alpha_0(a, m) \leq \epsilon_*(m), \quad (3.28)$$

where the constants $\alpha_0(a, m) \in]\Theta_0, 1[$ and $\epsilon_*(m) > 0$ are introduced in Theorem 3.1 and (3.16) respectively.

Theorem 3.3. (*General domains*)

Under the hypotheses (3.27) and (3.28), if Ω is not a disc, the following holds

$$\lim_{B \rightarrow \infty} \left(\sup_{|\alpha - \alpha_0(a, m)| \leq \epsilon_*(m)} |\partial_B \lambda_{1,\pm}(B, \alpha) - \beta(a, m, \alpha) - \alpha| \right) = 0. \quad (3.29)$$

Theorem 3.4. (*Disc domains*)

Assume that $\Omega = D(0, 1)$ is a disc. Given $a > 0$ and $m > m_*$, there exists a constant $B_0 > 1$ and a function $[B_0, \infty[\ni B \mapsto g(B)$ satisfying $\lim_{B \rightarrow \infty} g(B) = 0$, such that if $B \geq B_0$ and α satisfies (3.28), then

$$\partial_B \lambda_{1,+}(B, \alpha) \geq \alpha - \frac{1}{2}C_2(a, m, \alpha) + g(B). \quad (3.30)$$

In view of (3.26), we get as corollary of Theorems 3.3 and 3.4:

Theorem 3.5. Let $a > 0$ and $m > m_*$. Assume that $\Omega \subset \mathbb{R}^2$ has a smooth and compact boundary, and if Ω is a disc, assume in addition that,

$$\begin{aligned} \frac{1}{a} \left(a\alpha_0(a, m) + \frac{1}{2}C_2(a, m, \alpha_0(a, m)) \right) \left(1 - \frac{\Theta_0}{\alpha_0(a, m)} \right) \\ < \alpha_0(a, m) - \frac{1}{2}C_2(a, m, \alpha_0(a, m)). \end{aligned} \quad (3.31)$$

²Actually for z_0 small, $P[z, \alpha]$ is sectorial for all $z \in D_0$, its form domain is independent from z and the expression of the quadratic form associated to $P[z, \alpha]$ is analytic in z .

Then there exists $\kappa_0 > 0$ such that, for all $\kappa \geq \kappa_0$, there is a unique $H_*(\kappa) > 0$ solving the equation

$$\mu^{(1)}(\kappa, H_*(\kappa)) = 0,$$

where $\mu^{(1)}(\kappa, H)$ is introduced in (1.7).

Moreover, for all $H > H_*(\kappa)$ and $\kappa \geq \kappa_0$, $\mu^{(1)}(\kappa, H) > 0$.

Remark 3.6. Thanks to the asymptotic result of Theorem 3.1, the condition (3.31) is fulfilled for large values of m . Thus Theorem 3.1 and Theorem 3.5 imply Theorem 1.5.

Proof of Theorem 3.5. From [23], we get a constant κ_0 and a positive function $\delta(\kappa)$ such that $\lim_{\kappa \rightarrow \infty} \delta(\kappa) = 0$ and

$$\forall \kappa \in [\kappa_0, \infty[, \quad \mu^{(1)}(\kappa, H) = 0 \implies \left| H - \frac{\kappa}{\alpha_0(a, m)} \right| \leq \kappa \delta(\kappa).$$

It is moreover proved that one may find a solution $H > 0$ such that $\mu^{(1)}(\kappa, H) = 0$.

Let $H_*(\kappa) = \min\{H > 0 : \mu^{(1)}(\kappa, H) = 0\}$. It is sufficient to show that $\mu^{(1)}(\kappa, H) > 0$ for any magnetic field H satisfying

$$H_*(\kappa) < H \leq \frac{\kappa}{\alpha_0(a, m)} + \kappa \delta(\kappa). \quad (3.32)$$

Thus, for such a magnetic field, we put $B = B(\kappa; H) = \kappa H$ and $\alpha = \alpha(\kappa; H) = \kappa/H$ so that we can pick $\kappa_0 > 0$ sufficiently large such that the hypothesis (3.28) is valid, and hence the results of Theorems 3.3 and 3.4 hold.

Let us show that the function $H \mapsto \mu_1(\kappa, H)$ is strictly increasing in the interval

$$\left] \frac{\kappa}{\alpha_0(a, m)} - \kappa \delta(\kappa), \frac{\kappa}{\alpha_0(a, m)} + \kappa \delta(\kappa) \right[. \quad (3.33)$$

We assume that $\kappa \geq \kappa_0$. From (3.26), we write

$$\partial_H \mu_{\pm}^{(1)}(\kappa, H) = \kappa \left(\partial_B \lambda_{1, \pm}(B, \alpha) - \alpha \int_{\tilde{\Omega}} (V_a(x) + 1) |\phi_{B, \alpha}|^2 dx \right), \quad (3.34)$$

where $\phi_{B, \alpha}$ is an L^2 -normalized eigenfunction associated with the lowest eigenvalue $\mu_1(B, \alpha)$.

Assume first that Ω is not a disc. Invoking Theorem 3.3, we get, provided that κ_0 is large enough,

$$\partial_H \mu_{\pm}^{(1)}(\kappa, H) \geq \kappa \alpha \left(1 - \int_{\tilde{\Omega}} (V_a(x) + 1) |\phi_{B, \alpha}|^2 dx + o(1) \right), \quad (3.35)$$

where we use also that for H in the interval (3.33), $\beta(a, m, \alpha) = o(1)$ as $\kappa \rightarrow \infty$ (see Theorem 3.1).

Since $\phi_{B, \alpha}$ is normalized in $L^2(\tilde{\Omega})$ and an eigenfunction associated with $\mu_1(B, \alpha)$, we write,

$$\begin{aligned} & \alpha \left(1 - \int_{\tilde{\Omega}} (V_a(x) + 1) |\phi_{B, \alpha}|^2 dx \right) \\ &= \alpha \int_{\Omega} |\phi_{B, \alpha}|^2 dx - a \alpha \int_{\tilde{\Omega} \setminus \Omega} |\phi_{B, \alpha}|^2 dx \\ &= B^{-1} \left(\int_{\tilde{\Omega}} w_m(x) |(\nabla - iB\mathbf{F})\phi_{B, \alpha}|^2 dx - \mu_1(B, \alpha) \right). \end{aligned}$$

Using the min-max variational principle, we infer from the above,

$$\alpha \left(1 - \int_{\tilde{\Omega}} (V_a(x) + 1) |\phi_{B, \alpha}|^2 dx \right) \geq B^{-1} \left(\frac{1}{m} \mu^N(B) + \mu_1(B, \alpha) \right),$$

where $\mu^N(B)$ is the lowest eigenvalue of the Neumann magnetic Laplacian $-(\nabla - iB\mathbf{F})^2$ in $\tilde{\Omega}$.

It is a standard result that $\mu^N(B) = \Theta_0 B + o(B)$ as $B \rightarrow \infty$ (see [18]), and under the condition that H remains in the interval (3.33), it is proved in [23] that $\mu_1(B, \alpha) = o(B)$ as $B \rightarrow \infty$ (see Proposition 3.7 below). Thus, provided that H is in the interval (3.33) and κ_0 is large enough, we get

$$\alpha \left(1 - \int_{\tilde{\Omega}} (V_a(x) + 1) |\phi_{B,\alpha}|^2 dx \right) \geq \frac{\Theta_0}{m} + o(1),$$

and upon substitution in (3.35), we get when Ω is not a disc

$$\partial_H \mu_{\pm}^{(1)}(\kappa, H) > 0, \quad \forall \kappa \geq \kappa_0.$$

Now, we assume that Ω is a disc. In this case, Theorem 3.4 and (3.34) together yield,

$$\partial_H \mu_{\pm}^{(1)}(\kappa, H) \geq \kappa \left(\alpha - \frac{1}{2} C_2(a, m, \alpha) - \alpha(a+1) \int_{\tilde{\Omega} \setminus \Omega} |\phi_{B,\alpha}|^2 dx + o(1) \right). \quad (3.36)$$

Again, using $\int_{\tilde{\Omega}} |\phi_{B,\alpha}|^2 dx = 1$, we write,

$$\begin{aligned} & \alpha - \frac{1}{2} C_2(a, m, \alpha) - \alpha(a+1) \int_{\tilde{\Omega} \setminus \Omega} |\phi_{B,\alpha}|^2 dx \\ &= \left(\alpha - \frac{1}{2} C_2(a, m, \alpha) \right) \int_{\Omega} |\phi_{B,\alpha}|^2 dx - \left(a\alpha + \frac{1}{2} C_2(a, m, \alpha) \right) \int_{\tilde{\Omega} \setminus \Omega} |\phi_{B,\alpha}|^2 dx. \end{aligned}$$

Now, we use that $\phi_{B,\alpha}$ is an eigenfunction associated with $\mu_1(B, \alpha)$. From the identity $Q[B, \alpha](\phi_{B,\alpha}) = \mu_1(B, \alpha)$ and the min-max principle, we deduce,

$$a \int_{\tilde{\Omega} \setminus \Omega} |\phi_{B,\alpha}|^2 dx \leq \left(1 - \frac{\mu^N(B, \Omega)}{\alpha} \right) \int_{\Omega} |\phi_{B,\alpha}|^2 dx.$$

Here $\mu^N(B, \Omega)$ is the first eigenvalue of the Neumann magnetic Laplacian $-(\nabla - iB\mathbf{A})^2$ in Ω . Using again the asymptotic behavior of $\mu^N(B)$ as $B \rightarrow \infty$ and the condition that H remains in (3.33), we deduce,

$$a \int_{\tilde{\Omega} \setminus \Omega} |\phi_{B,\alpha}|^2 dx \leq \left(1 - \frac{\Theta_0}{\alpha_0(a, m)} + o(1) \right) \int_{\Omega} |\phi_{B,\alpha}|^2 dx.$$

Let us define,

$$\begin{aligned} \delta_0 &= \alpha - \frac{1}{2} C_2(a, m, \alpha_0(a, m)) \\ &\quad - \frac{1}{a} \left(a\alpha + \frac{1}{2} C_2(a, m, \alpha_0(a, m)) \right) \left(1 - \frac{\Theta_0}{\alpha_0(a, m)} \right). \end{aligned}$$

Under the condition (3.31), $\delta_0 > 0$. Under the hypothesis that H remains in the interval (3.33), we get provided that κ_0 is large enough,

$$\alpha - \frac{1}{2} C_2(a, m, \alpha) - \alpha(a+1) \int_{\tilde{\Omega} \setminus \Omega} |\phi_{B,\alpha}|^2 dx \geq \frac{\delta_0}{2},$$

for all $\kappa \geq \kappa_0$. Substituting in (3.36), we get the desired result that the function $H \mapsto \mu^{(1)}(\kappa, H)$ is strictly increasing in the interval (3.33) for all $\kappa \geq \kappa_0$. $\square$

To prove Theorem 3.3 we deal separately with the case where the domain Ω is a disc and the case where it is not. This will be the subject of the next sections, but we review first some of the known results concerning $\mu_1(B, \alpha)$.

3.3. Some facts concerning $\mu_1(B, \alpha)$. Let us recall the main results obtained in [23] for $\mu_1(B, \alpha)$, the bottom of the spectrum of the operator $P[B, \alpha]$ associated with the quadratic form (3.21). We mention that in [23] we deal with $\mu^{(1)}(\kappa, H)$ rather than $\mu_1(B, \alpha)$, but due to (3.24) we get equivalent results for $\mu_1(B, \alpha)$.

We start by giving the leading order term of $\mu_1(B, \alpha)$ as $B \rightarrow \infty$ ([23, Section 5]).

Proposition 3.7. *Given $a > 0$ and $m > 0$, there exist a constant $B_0 > 0$ and a function $[B_0, \infty[\ni B \mapsto g(B) \in \mathbb{R}_+$ satisfying $\lim_{B \rightarrow \infty} g(B) = 0$ such that, for any $\alpha > 0$ fulfilling (3.28), the following asymptotic holds:*

$$|\mu_1(B, \alpha) - \beta(a, m, \alpha)B| \leq g(B)B, \quad \forall B \geq B_0,$$

where $\beta(a, m, \alpha)$ is introduced in (3.6).

Letting $\phi_{B, \alpha}$ be a ground state of $P[B, \alpha]$, it is proved in [23, Section 6] that for $m > 1$, $\phi_{B, \alpha}$ is localized near the interior boundary $\partial\Omega$ in the sense precised in the next lemma.

Lemma 3.8. *Under the hypotheses of Proposition 3.7, given $a > 0$, $m > 1$ and an integer $k \in \mathbb{N}$, there exist positive constants B_k and C_k such that,*

$$\int_{\tilde{\Omega}} [\text{dist}(x, \partial\Omega)]^k |\phi_{B, \alpha}(x)|^2 dx \leq \frac{C_k}{\sqrt{B}^k} \|\phi_{B, \alpha}\|_{L^2(\tilde{\Omega})}^2, \quad (3.37)$$

$$\int_{\tilde{\Omega}} [\text{dist}(x, \partial\Omega)]^k |(\nabla - iB\mathbf{F})\phi_{B, \alpha}(x)|^2 dx \leq \frac{C_k}{\sqrt{B}^{k-2}} \|\phi_{B, \alpha}\|_{L^2(\tilde{\Omega})}^2, \quad (3.38)$$

for all $B \geq B_k$.

Under the additional hypotheses (3.27) and (3.28), i.e. α close to the constant $\alpha_0(a, m)$ and m strictly larger than the constant $m_* \geq 1$ introduced in (3.15), we get a two term asymptotic expansion for the eigenvalue $\mu_1(B, \alpha)$ (see [23, Propositions 7.1 and 7.3]).

Theorem 3.9. *Assume that $a > 0$ and m satisfies (3.27). There exist a constant $B_0 > 0$ and a function $[B_0, \infty[\mapsto g(B) \in \mathbb{R}_+$ satisfying $\lim_{B \rightarrow \infty} g(B) = 0$ such that, if α fulfills (3.28), then the following asymptotic holds:*

$$\left| \mu_1(B, \alpha) - \left(\beta(a, m, \alpha)B + C_1(a, m, \alpha)(\kappa_r)_{\max} \sqrt{B} \right) \right| \leq g(B) \sqrt{B}, \quad \forall B \geq B_0,$$

where $\beta(a, m, \alpha)$ is introduced in (3.6), $C_1(a, m, \alpha)$ the negative constant introduced in (3.12), κ_r denotes the scalar curvature of $\partial\Omega$ and $(\kappa_r)_{\max}$ denotes the maximum of κ_r .

It should be noticed that there exist negative constants c_1 and c_2 such that, for α satisfying (3.28),

$$c_1 \leq C_1(a, m, \alpha) \leq c_2 < 0.$$

3.4. Proof of Theorem 3.3: General domains. We assume in this section that the domain Ω is not a disc, still with smooth and compact boundary. Thus, the set

$$\Pi = \{x \in \partial\Omega : \kappa_r(x) = (\kappa_r)_{\max}\} \quad (3.39)$$

is not identical to $\partial\Omega$, i.e. $\Pi \neq \partial\Omega$.

In this case, we can deduce from the asymptotic expansion in Theorem 3.9 that any ground state $\phi_{B, \alpha}$ of $P[B, \alpha]$ is localized near the set Π . One weak version of expressing this rough statement is through the estimate of the following lemma, see [18, 20] for sharper statements.

Lemma 3.10. *Under the hypotheses of Theorem 3.9, for all $\epsilon_0 > 0$ and $N \in \mathbb{N}$, there exist constants $B_0 > 0$ and C_N such that,*

$$\int_{\{\text{dist}(x, \Pi) \geq \epsilon_0\}} |\phi_{B, \alpha}(x)|^2 dx \leq C_N B^{-N}, \quad \forall B \geq B_0.$$

Sketch of Proof of Lemma 3.10. We provide some details concerning the derivation of the estimate of Lemma 3.10. Using the decay estimates of Lemma 3.8, one can get the lower bound³,

$$Q[B, \alpha](\phi_{B, \alpha}) \geq \int_{\tilde{\Omega}} U_B(x) |\phi_{B, \alpha}(x)|^2 dx, \quad (3.40)$$

where

$$U_B(x) = \begin{cases} (\gamma - \alpha)B, & \text{if } \text{dist}(x, \partial\Omega) \geq 2B^{-1/6}, \\ \beta B + C_1 \kappa_r(p(x)) B^{1/2} - C_0 B^{1/3}, & \text{if } \text{dist}(x, \partial\Omega) \leq 2B^{-1/6}. \end{cases}$$

Here $\gamma \in]0, 1[$ is a given constant, $\beta = \beta(a, m, \alpha)$, $C_1 = C_1(a, m, \alpha)$, the constant $C_0 > 0$ depends on $a > 0$, $m > m_*$ and γ , and $p(x) \in \partial\Omega$ is defined by $|p(x) - x| = \text{dist}(x, \partial\Omega)$.

On the other hand, it follows from Theorem 3.9 that

$$Q[B, \alpha](\phi_{B, \alpha}) \leq \int_{\Omega} \left(\beta(a, m, \alpha)B + C_1(a, m, \alpha)(\kappa_r)_{\max} B^{1/2} + g(B) B^{1/2} \right) |\phi_{B, \alpha}|^2 dx. \quad (3.41)$$

Combining (3.40) and (3.41), one can prove the following exponential decay using standard (Agmon-type) techniques, (see [24, Theorem 6.5.4] for details)

$$\left\| \exp \left(\delta_0 B^{1/4} \text{dist}(x, \Pi) \right) \phi_{B, \alpha} \right\|_{L^2(\tilde{\Omega})} \leq M \|\phi_{B, \alpha}\|_{L^2(\tilde{\Omega})},$$

where $\delta_0 > 0$ and M are constants depending on a and m . Now the estimate of the lemma is just a consequence of the above exponential decay of $\phi_{B, \alpha}$. $\square$

We start now by following ideas of Fournais-Helffer in [9] to prove monotonicity of the eigenvalue

$$\lambda_1(B, \alpha) = \mu_1(B, \alpha) + \alpha B. \quad (3.42)$$

Notice that $\lambda_1(B, \alpha)$ is the eigenvalue of the operator

$$\tilde{P}[B, \alpha] = P[B, \alpha] + \alpha B$$

associated with the quadratic form

$$\phi \mapsto \tilde{Q}[B, \alpha](\phi) = Q[B, \alpha](\phi) + \alpha B \|\phi\|_{L^2(\tilde{\Omega})}^2, \quad (3.43)$$

where $Q[B, \alpha]$ is the quadratic form introduced in (3.21), and both operators $\tilde{P}[B, \alpha]$ and $P[B, \alpha]$ admit the same ground states. With this point of view, it is more convenient to adapt the proof of [9].

We proceed to prove Theorem 3.3.

³ This is a consequence of combining Proposition 5.3.3 and Theorem 4.3.8 in [24]. More details are given also in [24, pp. 179-180]

3.4.1. Gauge transformation. Using an idea from [9], we will apply a gauge transformation and work with a new magnetic potential $\mathbf{A}$ (instead of $\mathbf{F}$). In the new gauge, we will have that $|\mathbf{A}\phi_{B,\alpha}|$ is small in the L^2 -sense as $B \rightarrow \infty$. Notice that by Lemma 3.10 $\phi_{B,\alpha}$ is localized near the set Π , so it suffices to find a gauge where $\mathbf{A} = 0$ on Π .

To this end, we introduce adapted coordinates near the boundary of Ω . For a sufficiently small $t_0 > 0$, we introduce the open set

$$\Omega(t_0) = \{x \in \mathbb{R}^2 : \text{dist}(x, \partial\Omega) < t_0\}.$$

Let $s \mapsto \gamma(s)$ be the arc-length parametrization of $\partial\Omega$ and $\nu(s)$ the unit inward normal of $\partial\Omega$ at $\gamma(s)$.

When t_0 is sufficiently small, the transformation

$$\Phi : \left[-\frac{|\partial\Omega|}{2}, \frac{|\partial\Omega|}{2} \right] \times]-t_0, t_0[\ni (s, t) \mapsto \gamma(s) + t\nu(s) \in \Omega(t_0) \quad (3.44)$$

becomes a diffeomorphism whose Jacobian is $|D\Phi| = 1 - t\kappa_{\Gamma}(s)$. For $x \in \Omega(t_0)$, we put

$$\Phi^{-1}(x) = (s(x), t(x))$$

and we get in particular that

$$t(x) = \text{dist}(x, \partial\Omega) \quad \text{in } \Omega, \quad t(x) = -\text{dist}(x, \partial\Omega) \quad \text{outside } \Omega.$$

Just as in [9, Lemma 2.5], we get the following lemma.

Lemma 3.11. *Let $0 < \epsilon \leq \frac{1}{2} \min(t_0, |\partial\Omega|)$, $x_0 = \Phi(s_0, 0) \in \partial\Omega$ and*

$$\Omega(\epsilon, s_0) = \{x = \Phi(s, t) \in \tilde{\Omega} : |t| \leq \epsilon, |s - s_0| \geq \epsilon\}.$$

Then there exists a function $\varphi \in C_0^\infty(\mathbb{R}^2)$ such that $\mathbf{A} = \mathbf{F} + \nabla\varphi$ satisfies

$$|\mathbf{A}(x)| \leq C \text{dist}(x, \partial\Omega) \quad \text{in } \Omega(\epsilon, s_0),$$

where $C > 0$ depends only on Ω .

Proof of Theorem 3.3.

We have now all the prerequisites needed to apply the argument of Fournais-Helffer [9]. We include the details for the reader's convenience.

Recall that, for $\epsilon > 0$ sufficiently small and $z \in [B, B + \epsilon[$, we associate to $\lambda_1(z, \alpha)$ an analytic branch of eigenfunctions $z \mapsto \phi_{z,\alpha}$ such that

$$\tilde{P}[z, \alpha]\phi_{z,\alpha} = \lambda_1(z, \alpha)\phi_{z,\alpha} \quad \forall z \in [B, B + \epsilon[.$$

We may also in addition assume that $\|\phi_{z,\alpha}\|_{L^2(\tilde{\Omega})}^2 = 1$ for all $z \in [B, B + \epsilon[$.

Since Ω is not a disc, the set Π in (3.39) is different from Ω , and we can find $\epsilon_0 \in \frac{1}{2} \min(t_0, |\partial\Omega|)$ and $s_0 \in \partial\Omega$ such that

$$[s_0 - 2\epsilon_0, s_0 + 2\epsilon_0] \cap \Pi = \emptyset. \quad (3.45)$$

Let $\mathbf{A}$ and φ be respectively the vector field and the real-valued function defined in Lemma 3.11. Let $\hat{P}[z, \alpha]$ be the self adjoint operator associated to the quadratic form

$$H^1(\tilde{\Omega}) \ni u \mapsto \int_{\tilde{\Omega}} (w_m(x)|(\nabla - iz\mathbf{A})u|^2 + \alpha z(V_a(x) + 1)|u|^2) dx$$

where w_m and V_a are introduced in (1.15). Notice that the operators $\hat{P}[z, \alpha]$ and $\tilde{P}[z, \alpha]$ are unitary equivalent : $\hat{P}[z, \alpha]u = e^{iz\varphi}\tilde{P}[z, \alpha]e^{-iz\varphi}u$. Hence,

$$[B, B + \epsilon[\ni z \mapsto \hat{\phi}_{z,\alpha} := e^{iz\varphi}\phi_{z,\alpha}$$

is an analytic branch of eigenfunctions associated to the eigenvalue $\lambda_1(z, \alpha)$ of the operator $\hat{P}[z, \alpha]$.

Calculating,

$$\begin{aligned}\partial_B \lambda_{1,+}(B) &= \frac{d}{dz} \left(\widehat{Q}[B+z, \alpha](\widehat{\phi}_{z,\alpha}) \right) \Big|_{z=0_+} \\ &= \int_{\widetilde{\Omega}} 2w_m(x) \operatorname{Re} \left\langle -i\mathbf{A}\widehat{\phi}_{B,\alpha}, (\nabla - iB\mathbf{A})\widehat{\phi}_{B,\alpha} \right\rangle + \alpha (V_a(x) + \alpha) |\widehat{\phi}_{B,\alpha}(x)|^2 dx \\ &\quad + 2 \operatorname{Re} \widehat{Q}[B, \alpha] \left(\frac{d\widehat{\phi}_{B+z,\alpha}}{dz} \Big|_{z=0_+}, \widehat{\phi}_{B,\alpha} \right),\end{aligned}$$

where the last term on the r.h.s. above vanishes, since $\widehat{\phi}_{z,\alpha}$ is normalized with respect to the L^2 norm. Thus, for an arbitrary $\zeta \in \mathbb{R} \setminus \{0\}$, we may express $\partial_B \lambda_{1,+}(B)$ in the following way

$$\partial_B \lambda_{1,+}(B) = \frac{\widehat{Q}[B+\zeta, \alpha](\widehat{\phi}_{B,\alpha}) - \widehat{Q}[B, \alpha](\widehat{\phi}_{B,\alpha})}{\zeta} - \zeta \int_{\widetilde{\Omega}} |\mathbf{A}\widehat{\phi}_{B,\alpha}|^2 dx,$$

and invoking the min-max principle we get further when $\zeta > 0$

$$\partial_B \lambda_{1,+}(B) \geq \frac{\lambda_1(B+\zeta, \alpha) - \lambda_1(B, \alpha)}{\zeta} - \zeta \int_{\widetilde{\Omega}} |\mathbf{A}\phi_{B,\alpha}|^2 dx. \quad (3.46)$$

By Lemma 3.11, we may write,

$$\int_{\widetilde{\Omega}} |\mathbf{A}\phi_{B,\alpha}|^2 dx \leq C \int_{\widetilde{\Omega}} |\operatorname{dist}(x, \partial\Omega)|^2 |\phi_{B,\alpha}|^2 dx + \|\mathbf{A}\|_{L^\infty(\widetilde{\Omega})}^2 \int_{\widetilde{\Omega} \setminus \Omega(s_0, \epsilon_0)} |\phi_{B,\alpha}|^2 dx.$$

By our choice of s_0 and ϵ_0 , $\widetilde{\Omega} \setminus \Omega(s_0, \epsilon_0)$ is away from boundary points of maximal curvature. Thus, invoking Lemmas 3.37 and 3.10, we obtain a constant $B_0 > 0$ depending only on a and m such that

$$\int_{\widetilde{\Omega}} |\mathbf{A}\phi_{B,\alpha}|^2 dx \leq CB^{-1} \quad \text{for } B \geq B_0,$$

and we implement this last estimate in (3.46). Now, choosing $\zeta = \eta B$ in (3.46) with $\eta \in]0, 1[$ being arbitrary, we get from Proposition 3.7,

$$\begin{aligned}\partial_B \lambda_{1,+}(B) &\geq \beta(a, m, \alpha) + \alpha \\ &\quad - \frac{1+\eta}{\eta} g((1+\eta)B) - \frac{1}{\eta} g(B) - C\eta,\end{aligned} \quad (3.47)$$

where g is independent from α , and $g(B) \rightarrow 0$ as $B \rightarrow \infty$.

Applying the same argument to the left derivative $\partial_B \lambda_{1,-}(B)$, we get (the inequality being reversed since $z < 0$ in this case),

$$\begin{aligned}\partial_B \lambda_{1,-}(B) &\leq \beta(a, m, \alpha) + \alpha \\ &\quad + \frac{1+\eta}{\eta} g((1+\eta)B) + \frac{1}{\eta} g(B) + C\eta.\end{aligned} \quad (3.48)$$

Recall that analytic perturbation theory gives $\partial_B \lambda_{1,+}(B) \leq \partial_B \lambda_{1,-}(B)$ for all B . Therefore, (3.47) and (3.48) when combined together yield,

$$\limsup_{B \rightarrow \infty} \left(\sup_{|\alpha - \alpha_0(a, m)| \leq \epsilon_*(m)} |\partial_B \lambda_{1,\pm}(B, \alpha) - \beta(a, m, \alpha) - \alpha| \right) \leq C\eta.$$

Taking $\eta \rightarrow 0_+$ above, we get

$$\lim_{B \rightarrow \infty} \left(\sup_{|\alpha - \alpha_0(a, m)| \leq \epsilon_*(m)} |\partial_B \lambda_{1,\pm}(B, \alpha) - \beta(a, m, \alpha) - \alpha| \right) = 0.$$

□

3.5. Proof of Theorem 3.4: Disc domains. In order to handle disc domains, we need a refined asymptotic expansion of the first eigenvalue $\mu_1(B, \alpha)$ as $B \rightarrow \infty$. Let us introduce some notation. Given $a > 0$ and $m > m_*$, we introduce,

$$\delta(n, B) = n - \frac{1}{2}B - \xi(a, m, \alpha)\sqrt{B}. \quad (3.49)$$

Theorem 3.12. *Assume that $\Omega = D(0, 1)$. Given $a > 0$ and $m > m_*$, there exist a constant $B_0 > 1$ and a function $[B_0, \infty[\ni B \mapsto g(B)$ satisfying $\lim_{B \rightarrow \infty} g(B) = 0$, and if α satisfies (3.28), there exist real constants*

$$\delta_0 = \delta_0(a, m, \alpha), \quad \mathcal{C}_0 = \mathcal{C}_0(a, m, \alpha),$$

such that, for all $B \geq B_0$, the following expansion holds,

$$\left| \mu_1(B, \alpha) - \left(\beta(a, m, \alpha)B - C_1(a, m, \alpha)\sqrt{B} + C_2(a, m, \alpha)(\Delta_B^2 + \mathcal{C}_0) \right) \right| \leq g(B). \quad (3.50)$$

Here $C_2(a, m, \alpha)$ is introduced in (3.18) and

$$\Delta_B = \inf_{n \in \mathbb{Z}} |\delta(n, B) - \delta_0(a, m, \alpha)|.$$

The proof of Theorem 3.12 is very close to that of Theorem 2.5 in [8] and relies strongly on the fact that the eigenvalue (3.23) admits a non-degenerate minimum in ξ . For the convenience of the reader, we show in the appendix how the proof of [8] gives Theorem 3.12.

Recall the eigenvalue $\lambda_1(B, \alpha)$ introduced in (3.42). In view of the result of Theorem 3.12 above, Theorem 3.4 follows directly as [8, Theorem 2.5]. We omit thus the details.

4. EXPONENTIAL DECAY OF ORDER PARAMETERS

The objective of this section is to prove that, for $m > 1$, energy minimizing order parameters decay exponentially away from the boundary. This estimate will be useful for forthcoming works dedicated to finer properties of energy minimizers.

The main theorem is the following.

Theorem 4.1. *Assume that $a > 0$, $m > 1$ and let $b > 0$ be a given constant. There exist positive constants M , C , ε and κ_0 such that, if $(\psi, \mathbf{A})$ is a solution of (2.1) and the magnetic field verifies*

$$\frac{H}{\kappa} \geq 1 + b,$$

then

$$\int_{\tilde{\Omega}} e^{2\varepsilon\sqrt{\kappa H}t(x)} \left(|\psi|^2 + \frac{1}{\kappa H} |(\nabla - i\kappa H \mathbf{A})\psi|^2 \right) dx \leq C \int_{\{t(x) \leq \frac{M}{\kappa H}\}} |\psi|^2 dx. \quad (4.1)$$

Here $t(x) := \text{dist}(x, \partial\Omega)$.

When $m > m_*$ and the magnetic field $H = H_{C_3}(\kappa) - o(1)$, one should be able to prove a finer localization of ψ , near boundary points with maximal curvature. Actually, an estimate similar to Lemma 3.10 for the linear problem should also be valid for the solution ψ of the non-linear Ginzburg-Landau problem. However, to establish such an estimate will require a rather very technical work following previous papers [20, 27], so that we do not carry it out. Let us only notice here that—just as for the linear case—such a localisation estimate is essentially due to the asymptotic expansion of the first eigenvalue stated in Theorem 3.9.

Proof of Theorem 4.1.

If $\psi \equiv 0$, then the estimate of the theorem is evidently true. Thus, thanks to Theorem 2.3, we may restrict ourselves to magnetic fields H satisfying

$$H \leq C\kappa$$

for some sufficiently large positive constant C .

Let $\tilde{\chi} \in C^\infty(\mathbb{R})$ be a non-decreasing function with

$$\tilde{\chi} = 1, \quad \text{on } [1, \infty), \quad \tilde{\chi} = 0, \quad \text{on } (-\infty, 1/2]. \quad (4.2)$$

Define χ on $\tilde{\Omega}$ by $\chi(x) = \tilde{\chi}(\sqrt{\kappa H}t(x))$. Define furthermore, the weighted localisation function f by

$$f(x) = \chi(x) \exp(\varepsilon \sqrt{\kappa H}t(x)) \quad (4.3)$$

We calculate, using (2.1)

$$\begin{aligned} & \int_{\tilde{\Omega}} w_m(x) \left(|(\nabla - i\kappa H \mathbf{A})f\psi|^2 - |\nabla f|^2 |\psi|^2 \right) dx + a\kappa^2 \int_{\tilde{\Omega} \setminus \Omega} |f\psi|^2 dx \\ &= \kappa^2 \int_{\tilde{\Omega}} (|\psi|^2 - |\psi|^4) f^2 dx \\ &\leq \kappa^2 \int_{\Omega} f^2 |\psi|^2 dx \end{aligned} \quad (4.4)$$

Now, using Lemma 4.2 below, we can estimate

$$\begin{aligned} & \int_{\tilde{\Omega}} w_m(x) |(\nabla - i\kappa H \mathbf{A})f\psi|^2 dx \\ &= \int_{\Omega} |(\nabla - i\kappa H \mathbf{A})f\psi|^2 dx + m^{-1} \int_{\tilde{\Omega} \setminus \Omega} |(\nabla - i\kappa H \mathbf{A})f\psi|^2 dx \\ &\geq \kappa H (1 - C/\sqrt{\kappa H}) \int_{\Omega} |f\psi|^2 dx. \end{aligned} \quad (4.5)$$

Combining (4.4) and (4.5) we find

$$\left(\kappa H \frac{b}{1+b} - C\sqrt{\kappa H} \right) \int_{\Omega} |f\psi|^2 dx + a\kappa^2 \int_{\tilde{\Omega}} |f\psi|^2 dx \leq \int_{\tilde{\Omega}} |\nabla f|^2 |\psi|^2 dx. \quad (4.6)$$

We estimate the last term

$$\int_{\tilde{\Omega}} |\nabla f|^2 |\psi|^2 dx \leq 2\varepsilon^2 \kappa H \int_{\tilde{\Omega}} |f\psi|^2 dx + C\kappa H \int_{\{\sqrt{\kappa H}t(x) \leq 1\}} |\psi(x)|^2 dx. \quad (4.7)$$

Therefore we get, choosing ε sufficiently small and for κH sufficiently large,

$$\int_{\tilde{\Omega}} |f\psi|^2 dx \leq C \int_{\{\sqrt{\kappa H}t(x) \leq 1\}} |\psi(x)|^2 dx. \quad (4.8)$$

This implies the weighted L^2 -bound in (4.1),

$$\int_{\tilde{\Omega}} e^{2\varepsilon \sqrt{\kappa H}t} |\psi(x)|^2 dx \leq C \int_{\{\sqrt{\kappa H}t(x) \leq 1\}} |\psi(x)|^2 dx. \quad (4.9)$$

Inserting (4.9) in (4.4) (and using the same considerations) yields the weighted bound on $(\nabla - i\kappa H \mathbf{A})\psi$. $\square$

Lemma 4.2. *There exist constants C_0, C_1 such that if $(\psi, \mathbf{A})$ is a solution of (2.1) with $\kappa(H - \kappa) \geq C_0$, then*

$$\|(\nabla - i\kappa H \mathbf{A})\phi\|_{L^2(\Omega)}^2 \geq \kappa H \left(1 - \frac{C_1}{\sqrt{H(H - \kappa)}}\right) \|\phi\|_{L^2(\Omega)}^2, \quad (4.10)$$

for all $\phi \in C_0^\infty(\Omega)$. Also,

$$\|(\nabla - i\kappa H \mathbf{A})\phi\|_{L^2(\tilde{\Omega} \setminus \Omega)}^2 \geq \kappa H \left(1 - \frac{C_1}{\sqrt{H(H - \kappa)}}\right) \|\phi\|_{L^2(\tilde{\Omega} \setminus \Omega)}^2, \quad (4.11)$$

for all $\phi \in C_0^\infty(\tilde{\Omega} \setminus \Omega)$.

Proof. We only prove (4.10) the proof of (4.11) being identical.

We estimate, using the compact support of ϕ and the standard magnetic estimate from [2, Thm 2.9] (or [7, Lemma 2.4.1])

$$\begin{aligned} \|(\nabla - i\kappa H \mathbf{A})\phi\|_{L^2(\Omega)}^2 &\geq \kappa H \int_{\Omega} |\operatorname{curl} \mathbf{A}|^2 dx \\ &\geq \kappa H \|\phi\|_2^2 - (\kappa H) \|\operatorname{curl} \mathbf{A} - 1\|_2 \|\phi\|_4^2. \end{aligned} \quad (4.12)$$

By Lemma 2.2 and the weak decay estimate of Lemma 2.6, we have

$$\|\operatorname{curl} \mathbf{A} - 1\|_2 \leq \frac{C}{H \sqrt{\kappa(H - \kappa)}}. \quad (4.13)$$

Furthermore, by the Sobolev inequality and scaling followed by the diamagnetic inequality, we find

$$\begin{aligned} \|\phi\|_4^2 &\leq C_{\text{Sob}} \left(\eta \|\nabla |\phi|\|_2^2 + \eta^{-1} \|\phi\|_2^2 \right) \\ &\leq C_{\text{Sob}} \left(\eta \|(\nabla - i\kappa H \mathbf{A})\phi\|_2^2 + \eta^{-1} \|\phi\|_2^2 \right), \end{aligned} \quad (4.14)$$

where C_{Sob} is a universal constant and $\eta > 0$ is a parameter that we can choose freely. We make the choice $\eta = 1/\sqrt{\kappa H}$. Combining (4.12), (4.13) and (4.14) yields

$$\left(1 + \frac{C}{\sqrt{H(H - \kappa)}}\right) \|(\nabla - i\kappa H \mathbf{A})\phi\|_{L^2(\Omega)}^2 \geq \kappa H \left(1 - \frac{C}{\sqrt{H(H - \kappa)}}\right) \|\phi\|_{L^2(\Omega)}^2, \quad (4.15)$$

from which (4.10) follows. $\square$

ACKNOWLEDGEMENTS

The authors were supported by the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement n° 202859. SF is also supported by the Danish Research Council and the Lundbeck Foundation.

APPENDIX A. IMPROVED EIGENVALUE ESTIMATE FOR THE DISC

The aim of this appendix is to prove Theorem 3.12. By assumption, $\Omega = D(0, 1)$ and $D(0, 1 + r) \subset \tilde{\Omega}$ for some $r > 0$.

Let $D(t) = \{x \in \mathbb{R}^2 : |x| \leq t\}$ be the disc with radius t . Let $\tilde{Q}_B$ be the quadratic form

$$\tilde{Q}_B[u] = \int_{D(1+r) \setminus D(\frac{1}{2})} \left(w_m(x) |(\nabla - iB\mathbf{F})u|^2 + \alpha V_a(x) |u|^2 \right) dx,$$

with domain $\{u \in H^1(D(1+r) \setminus D(\frac{1}{2})) \mid u(x) = 0 \text{ on } |x| = \frac{1}{2} \text{ and } |x| = 1+r\}$. Here w_m and V_a are as in (1.15), and we emphasize that, for the sake of simplicity, we omit the dependence on a , m and α from the notation.

Let $\tilde{\mu}_1(B, \alpha)$ be the lowest eigenvalue of the corresponding self-adjoint operator. Using the variational principle and the decay of the ground state (Lemma 3.8), we see that,

$$\mu_1(B, \alpha) = \tilde{\mu}_1(B, \alpha) + \mathcal{O}(B^{-\infty}). \quad (\text{A.1})$$

So, it is sufficient to prove (3.50) with $\mu_1(B, \alpha)$ replaced by $\tilde{\mu}_1(B, \alpha)$.

By changing to boundary coordinates (s, t) (defined by (3.44)) the quadratic form $\tilde{Q}_B[u]$ becomes,

$$\tilde{Q}_B[u] = \int_0^{2\pi} \int_{-r}^{1/2} \tilde{w}_m(t) \left[(1-t)^{-2} |(D_s - B\tilde{A}_1)u|^2 + |D_t u|^2 \right] (1-t) dt ds \quad (\text{A.2})$$

$$+ \int_0^{2\pi} \int_{-r}^{1/2} \alpha \tilde{V}_a(t) (1-t) |u|^2 dt ds, \quad (\text{A.3})$$

$$\|u\|_{L^2}^2 = \int_0^{2\pi} \int_{-r}^{1/2} (1-t) |u|^2 dt ds, \quad \tilde{A}_1 = \frac{1}{2} - t + \frac{t^2}{2}.$$

Here

$$\tilde{w}_m(t) = \begin{cases} 1, & \text{if } t > 0 \\ \frac{1}{m}, & \text{if } t < 0 \end{cases} \quad \tilde{V}_a(t) = \begin{cases} -1, & \text{if } t > 0 \\ a, & \text{if } t < 0. \end{cases} \quad (\text{A.4})$$

Performing the scaling $\tau = \sqrt{B}t$ and decomposing in Fourier modes, we find

$$\tilde{\mu}_1(B, \alpha) = B \inf_{n \in \mathbb{Z}} e_{\delta(n, B), B}. \quad (\text{A.5})$$

Here the function $\delta(m, B)$ was defined in (3.49) and $e_{\delta, B}$ is the lowest eigenvalue of the quadratic form $q_{\delta, B}$ on $L^2((-\sqrt{B}r, \sqrt{B}/2); (1 - \sqrt{B}\tau)d\tau)$ (with Dirichlet condition, $u(\tau) = 0$, at $\tau = -\sqrt{B}r$ and $\tau = \sqrt{B}/2$),

$$\begin{aligned} q_{\delta, B}[\phi] = & \int_{-\sqrt{B}r}^{\sqrt{B}/2} \tilde{w}_m(\tau) \left[\left(1 - \frac{\tau}{\sqrt{B}}\right)^{-1} \left((\tau - \xi) + B^{-\frac{1}{2}}\left(\delta - \frac{\tau^2}{2}\right)\right)^2 + \left(1 - \frac{\tau}{\sqrt{B}}\right) |\phi'(\tau)|^2 \right] d\tau \\ & + \alpha \int_{-\sqrt{B}r}^{\sqrt{B}/2} \left(1 - \frac{\tau}{\sqrt{B}}\right) \tilde{V}_a(t) |\phi(\tau)|^2 d\tau. \end{aligned} \quad (\text{A.6})$$

Here, $\xi = \xi(a, m, \alpha)$ by convention (this makes sense provided that $m > m_*$, and $\alpha \in [\alpha_0(a, m) - \epsilon_*(m), \alpha_0(a, m) + \epsilon_*(m)]$). We will only consider δ varying in a fixed bounded set. This is justified since it follows from [23, Proposition 4.6] that for all $C > 0$ there exists $D > 0$ such that if $|\delta| > D$ and $B > D$, then

$$e_{\delta, B} \geq \beta(a, m, \alpha) + C_1(a, m, \alpha) B^{-\frac{1}{2}} + CB^{-1}.$$

Furthermore, for δ varying in a fixed bounded set, we know (from the analysis of the operator (3.4), especially that the minimum of (3.5) in ξ is non-degenerate) that there exists a $d > 0$ such that if $B > d^{-1}$, then the spectrum of $q_{\delta, B}$ contained in $]-\infty, \beta(a, m, \alpha) + d[$ consists of exactly one simple eigenvalue.

The self-adjoint operator $\mathfrak{h}(\delta, B)$ associated to $q_{\delta, B}$ (on the space $L^2((-\sqrt{B}r, \sqrt{B}/2); (1 - \sqrt{B}\tau)d\tau)$) is

$$\begin{aligned} \mathfrak{h}(\delta, B) = & -\left(1 - \frac{\tau}{\sqrt{B}}\right)^{-1} \frac{d}{d\tau} \tilde{w}_m(\tau) \left(1 - \frac{\tau}{\sqrt{B}}\right) \frac{d}{d\tau} \\ & + \tilde{w}_m(\tau) \left[\left(1 - \frac{\tau}{\sqrt{B}}\right)^{-2} \left((\tau - \xi) + B^{-\frac{1}{2}}\left(\delta - \frac{\tau^2}{2}\right)\right)^2 \right] + \alpha \tilde{V}_a(\tau). \end{aligned} \quad (\text{A.7})$$

We will write down an explicit test function for $\mathfrak{h}(\delta, B)$ in (A.12) below, giving $e_{\delta, B}$ up to an error of order $\mathcal{O}(B^{-\frac{3}{2}})$ (locally uniformly in δ).

We can formally develop $\mathfrak{h}(\delta, B)$ as

$$\mathfrak{h}(\delta, B) = \mathfrak{h}_0 + B^{-\frac{1}{2}} \mathfrak{h}_1 + B^{-1} \mathfrak{h}_2 + \mathcal{O}(B^{-\frac{3}{2}}).$$

with

$$\begin{aligned} \mathfrak{h}_0 &= -\frac{d}{d\tau}\tilde{w}_m(\tau)\frac{d}{d\tau} + \tilde{w}_m(\tau)(\tau - \xi)^2 + \alpha\tilde{V}_a(\tau) \quad (= H[a, m, \alpha; \xi]) , \\ \mathfrak{h}_1 &= \tilde{w}_m(\tau) \left[\frac{d}{d\tau} + 2(\tau - \xi)(\delta - \frac{\tau^2}{2}) + 2\tau(\tau - \xi)^2 \right] , \\ \mathfrak{h}_2 &= \tilde{w}_m(\tau) \left[\tau\frac{d}{d\tau} + (\delta - \frac{\tau^2}{2})^2 + 4\tau(\tau - \xi)(\delta - \frac{\tau^2}{2}) + 3\tau^2(\tau - \xi)^2 \right] . \end{aligned} \quad (\text{A.8})$$

Let u_0 be the known ground state eigenfunction of $H[a, m, \alpha; \xi]$ with eigenvalue $\beta(a, m, \alpha)$. Here, by $H[a, m, \alpha; \xi]$, we mean the operator (3.4) with $\xi = \xi(a, m, \alpha)$, considered as a self-adjoint operator on $L^2(\mathbb{R}; d\tau)$. For ease of notation we will write $\mathfrak{h}_0$ instead of $H[a, m, \alpha; \xi]$, since they are the same formal differential operators. Let R_0 be the regularized resolvent of $\mathfrak{h}_0$, which is defined by

$$R_0\phi = \begin{cases} (\mathfrak{h}_0 - \beta(a, m, \alpha))^{-1}\phi , & \int \phi(\tau)u_0(\tau) d\tau = 0 , \\ 0 , & \phi \parallel u_0 . \end{cases}$$

Let λ_1 and λ_2 be given by

$$\begin{aligned} \lambda_1 &:= \langle u_0 | \mathfrak{h}_1 u_0 \rangle_{L^2(\mathbb{R}; d\tau)} , \\ \lambda_2 &:= \lambda_{2,1} + \lambda_{2,2} , \\ \lambda_{2,1} &:= \langle u_0 | \mathfrak{h}_2 u_0 \rangle_{L^2(\mathbb{R}; d\tau)} , \quad \lambda_{2,2} := \langle u_0 | (\mathfrak{h}_1 - \lambda_1)u_1 \rangle_{L^2(\mathbb{R}; d\tau)} , \end{aligned} \quad (\text{A.9})$$

The functions u_1, u_2 are given as

$$u_1 = -R_0(\mathfrak{h}_1 - \lambda_1)u_0 , \quad u_2 = -R_0\{(\mathfrak{h}_1 - \lambda_1)u_1 + (\mathfrak{h}_2 - \lambda_2)u_0\} . \quad (\text{A.10})$$

Using the same type of argument in [21, Proposition II.10] or [10, Lemma A.5], we can prove that R_0 preserves exponential decay at infinity, i.e. $u_0(t)$, $u_1(t)$, $u_2(t)$ and their derivatives decay exponentially fast as $|t| \rightarrow \infty$.

Let $\chi \in C_0^\infty(\mathbb{R})$ be a usual cut-off function, such that

$$\chi(t) = 1 \quad \text{for } |t| \leq \frac{1}{8} , \quad \text{supp } \chi \subset [-\frac{1}{4}, \frac{1}{4}] , \quad (\text{A.11})$$

and let $\chi_B(\tau) = \chi(\tau B^{-\frac{1}{4}})$.

We define the following trial state,

$$\psi := \chi_B\{u_0 + B^{-\frac{1}{2}}u_1 + B^{-1}u_2\} . \quad (\text{A.12})$$

Using the exponential decay of the involved functions, we get after a calculation,

$$\begin{aligned} \|\{\mathfrak{h}(\delta, B) - (\beta(a, m, \alpha) + \lambda_1 B^{-\frac{1}{2}} + \lambda_2 B^{-1})\}\psi\|_{L^2([- \sqrt{B}/2, \sqrt{B}/2]; (1 - \sqrt{B}\tau)d\tau)} \\ = \mathcal{O}(B^{-\frac{3}{2}}) , \end{aligned} \quad (\text{A.13})$$

$$\|\psi\|_{L^2([- \sqrt{B}/2, \sqrt{B}/2]; (1 - \sqrt{B}\tau)d\tau)} = 1 + \mathcal{O}(B^{-\frac{1}{2}}) , \quad (\text{A.14})$$

where the constant in $\mathcal{O}$ is uniform for δ in bounded sets. Applying the spectral theorem, and noticing that $\beta(a, m, \alpha)$ is an isolated eigenvalue for the operator $\mathfrak{h}_0$, we deduce that (uniformly for δ varying in bounded sets),

$$e_{\delta, B} = \beta(a, m, \alpha) + \lambda_1 B^{-\frac{1}{2}} + \lambda_2 B^{-1} + \mathcal{O}(B^{-\frac{3}{2}}) . \quad (\text{A.15})$$

It remains to calculate λ_1, λ_2 and, in particular, deduce their dependence on δ .

Writing,

$$2(\tau - \xi) \left(\delta - \frac{\tau^2}{2} \right) + 2\tau(\tau - \xi)^2 = (\tau - \xi)^3 - (\xi^2 + 2\delta)(\tau - \xi) ,$$

and using

$$\int_{-\infty}^{\infty} \tilde{w}_m(\tau)(\tau - \xi)u_0^2 d\tau = 0 ,$$

we get,

$$\lambda_1 = C_1(a, m, \alpha),$$

where $C_1(a, m, \alpha)$ is introduced in (3.9). In particular, λ_1 is independent of δ .

We do not need to calculate λ_2 explicitly. Notice that $\lambda_2(\delta)$ is a quadratic polynomial as a function of δ . We find the coefficient to δ^2 as equal to,

$$\int_{\mathbb{R}} \tilde{w}_m(\tau) |u_0(\tau)|^2 d\tau - 4I_2,$$

with

$$I_2 := \langle u_0, \tilde{w}_m(\tau)(\tau - \xi) R_0 \tilde{w}_m(\tau - \xi) u_0 \rangle. \quad (\text{A.16})$$

Therefore, there exist constants $\delta_0, C_0 \in \mathbb{R}$ such that

$$\lambda_2 = \left(\int_{\mathbb{R}} \tilde{w}_m(\tau) |u_0(\tau)|^2 d\tau - 4I_2 \right) ((\delta - \delta_0)^2 + C_0).$$

Recalling the definition of the constant $C_2(a, m, \alpha)$ in (3.18), the above formula becomes,

$$\lambda_2 = C_2(a, m, \alpha) ((\delta - \delta_0)^2 + C_0).$$

In light of (A.1) and (A.5), we need only to show that $C_2(a, m, \alpha) > 0$ to finish the proof Theorem 3.12.

Notice that we work under the hypothesis $m \geq m_*$. This implies that ξ is the unique, non-degenerate minimum of the function (see (3.5) and (3.15))

$$z \mapsto \mu(z) := \mu_1(a, m, \alpha; z).$$

In particular,

$$\mu''(\xi) > 0.$$

Now, exactly as shown in [9, Proposition A.3], it holds that

$$C_2(a, m, \alpha) = \frac{1}{2} \mu''(\xi), \quad (\text{A.17})$$

yielding thus the desired property regarding the sign of $C_2(a, m, \alpha)$. This finishes the proof of Theorem 3.12. $\square$

For the sake of the reader's convenience, we include some details concerning the derivation of (A.17).

Sketch of the proof of (A.17). Let us introduce,

$$E(z) = \mu(z + \xi), \quad H(z) = -\frac{d}{d\tau} \tilde{w}_m \frac{d}{d\tau} + \tilde{w}_m(t)(t - \xi - z)^2 + \alpha V_a(t),$$

together with an analytic family of eigenfunctions $z \mapsto \phi(z) \in L^2(\mathbb{R})$ such that

$$\|\phi(z)\|_{L^2(\mathbb{R})}^2 = 1, \quad H(z)\phi(z) = E(z)\phi(z), \quad \phi(0) = u_0.$$

By differentiating the identity $\|\phi(z)\|^2 = 1$ twice with respect to z , we get

$$2\operatorname{Re}\langle \phi'(0), u_0 \rangle = 0, \quad \operatorname{Re}\langle \phi''(0), u_0 \rangle = -\|\phi'(0)\|^2. \quad (\text{A.18})$$

Differentiating the relation $H(z)\phi(z) = E(z)\phi(z)$ we get since $E(z)$ is minimal for $z = 0$,

$$(H(0) - E(0))\phi'(0) = 2\tilde{w}_m(t)(t - \xi)u_0.$$

Since the functions $\tilde{w}_m(t - \xi)u_0$ and u_0 are orthogonal in $L^2(\mathbb{R})$, we get

$$\phi'(0) = 2(H(0) - E(0))^{-1}(\tilde{w}_m(t - \xi)u_0) + c u_0, \quad (\text{A.19})$$

for some constant $c \in i\mathbb{R}$.

Differentiating twice the relation $E(z) = \langle \phi(z), H(z)\phi(z) \rangle$, we get,

$$E''(0) = 2E(0)\operatorname{Re}\langle \phi'(0), u_0 \rangle - 8\operatorname{Re}\langle \phi'(0), \tilde{w}_m(t - \xi)u_0 \rangle \\ + 2\operatorname{Re}\langle \phi'(0), H(0)\phi'(0) \rangle + 2\langle u_0, \tilde{w}_m u_0 \rangle. \quad (\text{A.20})$$

Substituting (A.18) and (A.19) in (A.20), we get,

$$E''(0) = 2 \left(\int_R \tilde{w}_m(\tau) |u_0(\tau)|^2 d\tau - 4I_2 \right)$$

with I_2 introduced in (A.16). Recalling the definition of the constant $C_2(a, m, \alpha)$ we get the desired relation in (A.17). $\square$

REFERENCES

- [1] S. AGMON, A. DOUGLIS, L. NIRENBERG. Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I. *Comm. Pure Appl. Math.* **12** 623-727 (1959).
- [2] J. AVRON, I. HERBST, AND B. SIMON. Schrödinger operators with magnetic fields I. General Interactions. *Duke Math. J.* **45** 847-883 (1978).
- [3] V. BONNAILLIE-NOËL, S. FOURNAIS. Superconductivity in domains with corners. *Rev. Math. Phys.* **19** (6) 607-637 (2007).
- [4] S.J. CHAPMAN, Q. DU, M.D. GUNZBURGER, A Ginzburg Landau type model of superconducting/normal junctions including Josephson junctions, *European J. Appl. Math.* **6** (2) 97-114 (1996).
- [5] Q. DU, M. GUNZBURGER, J. PETERSON. Analysis and approximation of the Ginzburg-Landau model of superconductivity. *SIAM Rev.* **34** 45-81 (1992).
- [6] L.C. EVANS. *Partial Differential Equations*. Graduate Studies in Mathematics, 19. American Mathematical Society, Providence, RI, 1998.
- [7] S. FOURNAIS, B. HELFFER. *Spectral Methods in Surface Superconductivity*. Monograph submitted.
- [8] S. FOURNAIS, B. HELFFER. On the third critical field in Ginzburg-Landau theory. *Comm. Math. Phys.* **226** (1) 153-196 (2006).
- [9] S. FOURNAIS, B. HELFFER. Strong diamagnetism in general domains and applications. *Ann. Inst. Fourier.* **57** (7) 2389-2400 (2007).
- [10] S. FOURNAIS, B. HELFFER. Accurate eigenvalue asymptotics for the magnetic Neumann Laplacian. *Ann. Inst. Fourier.* (2006).
- [11] S. FOURNAIS, B. HELFFER. On the Ginzburg-Landau critical field in three dimensions. *Commun. Pure Appl. Math.* **62** 215-241 (2009).
- [12] S. FOURNAIS, A. KACHMAR. Nucleation of bulk superconductivity close to critical magnetic field. In preparation.
- [13] P.G. DE GENNES, *Superconductivity of metals and alloys*, Benjamin (1966).
- [14] P.G. DE GENNES, Boundary effects in superconductors, *Rev. Mod. Phys.* January 1964.
- [15] P.G. DE GENNES, J.P. HURAU, Proximity effects under magnetic fields II- Interpretation of 'breakdown', *Phys. Lett.* **17** (3) 181-182 (1965).
- [16] T. GIORGI, Superconductors surrounded by normal materials, *Proc. Roy. Soc. Edinburgh*, **135A** 331-356 (2005).
- [17] T. GIORGI AND D. PHILLIPS. The breakdown of superconductivity due to strong fields for the Ginzburg-Landau model. *SIAM J. Math. Anal.* **30** (1999), p. 341-359.
- [18] B. HELFFER, A. MORAME. Magnetic bottles in connection with superconductivity. *J. Funct. Anal.* (2) 604-680 (2001).
- [19] B. HELFFER, A. MORAME. Magnetic bottles for the Neumann problem: curvature effect in the case of dimension 3 (general case). *Ann. Ec. Norm. Sup.* **37** (2004) p. 105-170.
- [20] B. HELFFER, X.B. PAN. Upper critical field and location of surface nucleation of superconductivity. *Ann. Inst. H. Poincaré, analyse non-linéaire* **20** (1) 145-181 (2003).
- [21] A. KACHMAR. On the ground state energy for a magnetic Schrödinger operator and the effect of the DeGennes boundary condition. *J. Math. Phys.* **47** 072106 (32 pp.) (2006).
- [22] A. KACHMAR. On the perfect superconducting solution for a generalized Ginzburg-Landau equation. *Asymptot. Anal.* **54** (3-4) (2007).
- [23] A. KACHMAR. On the stability of normal states for a generalized Ginzburg-Landau model. *Asymptot. Anal.* **55** (3-4) 145-201 (2007).

- [24] A. KACHMAR. *Problèmes aux limites issus de la supraconductivité, estimations semi-classiques et comportement asymptotique des solutions*. Ph.D. Thesis, Université Paris-Sud (2007).
- [25] T. KATO. *Perturbation Theory for Linear Operators*. Berlin: Springer-Verlag, 1976.
- [26] K. LU, X.B. PAN. Surface nucleation of superconductivity in 3-dimensions. *J. Differential Equations* **168** no. 2 (2000), 386-452.
- [27] K. LU, X.B. PAN. Estimates of the upper critical field of the Ginzburg-Landau equations of superconductivity. *Physica D* **127** no. 1-2, 73-104 (1999).
- [28] X.B. PAN. Surface superconductivity in applied magnetic fields above H_{c2} . *Comm. Math. Phys.* **228** (2) 327-370 (2002).
- [29] E. SANDIER, S. SERFATY. *Vortices in the Magnetic Ginzburg-Landau Model*. Progress in Nonlinear Differential Equations and Applications, Vol. 70, Birkhauser, 2007.
- [30] S. SERFATY. Local minimizers for the Ginzburg-Landau energy near critical magnetic field. Part I. *Commun. Contemp. Math.* **1** (2) 213-254 (1999).
- [31] S. SERFATY. Stable Configurations in Superconductivity : Uniqueness, Multiplicity and Vortex-Nucleation. *Archive for Rational Mechanics and Analysis*, **149** 329-365 (1999).
- [32] R. TEMAM. *Navier-Stokes Equation. Theory and Numerical Analysis*. (Revised version) Elsevier Science Publishers B.V. Amsterdam, 1984.

(S. FOURNAIS AND A. KACHMAR) DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF AARHUS, NY MUNKEGADE, BUILDING 1530, DK-8000 ÅRHUS C
E-mail address, S. Fournais: `fournais@imf.au.dk`
E-mail address, A. Kachmar: `ayman.kachmar@math.u-psud.fr`